

<p>Deductive Reasoning in the Lifelong Continuing Education of a Cardiovascular Surgeon Frank C. Spencer, MD, New York</p>	1177
<p>Prevention of Paraplegia Associated With Resection of Extensive Thoracic Aneurysms Akio Wakabayashi, MD, John E. Connolly, MD, Irvine, Calif Bypass and preservation of posterior aortic wall below D-9 are safeguards.</p>	1186
<p>Vascular Complications of Intra-aortic Balloon Pumping Joseph Alpert, MD; Ezhuthachan K. Bhaktan, MD; Isaac Gielchinsky, MD; Lawrence Gilbert, MD; Bruce J. Brener, MD; Donald K. Brief, MD; Victor Parsonnet, MD, Newark, NJ Limb ischemia more frequent than believe.</p>	1190
<p>Selective Intracavitary and Coronary Hypothermic Cardioplegia for Myocardial Preservation: Clinical, Physiologic, and Ultrastructural Evaluation Arie Schachner, MD; George Schimert, MD; Thomas Z. Lajos, MD; Arthur H. Lee, MD; Mario Montes, MD; Anand Chaudhry, PhD; Peter Schaefer, PhD; Adrian Vladutin, MD; John H. Siegel, MD, Buffalo, NY Less intraoperative myocardial damage.</p>	1197
<p>Results of Reoperation for Failed Aortocoronary Bypass Grafts Clarence S. Thomas, Jr, MD; William C. Alford, Jr, MD; George R. Burrus, MD; Robert A. Frist, MD; William S. Stoney, MD, Nashville, Tenn No increased risk but worse long term results than initial procedures.</p>	1210
<p>Myocardial Revascularization After Acute Myocardial Infarction Hooshang Bolooki, MD, FRCS(C), Abelardo Vargas, MD, Miami Four hours postinfarct maximum interval for reliable revascularization.</p>	1216
<p>Prolonged Pulseless Perfusion in Unanesthetized Calves G. Gilbert Johnston, MD; Frederick Hammill, MD; Ulla Marzec; Dava Gerard, MD; Kaj Johansen, MD; Ralph B. Dilley, MD; Eugene F. Bernstein, MD, PhD, La Jolla, Calif No untoward effects from nonpulsatile bypass for up to 14 days.</p>	1225

Contents continued on p 1172.

Official Publication, International Cardiovascular Society and Western Surgical Association

EDITORIAL BOARD

Richard Warren, MD, Boston
Chief Editor

Robert M. Goldwyn, MD, Brookline, Mass
Assistant Chief Editor

Arthur E. Baue, MD, New Haven, Conn

Jerome J. DeCosse, MD, Milwaukee

Eric W. Fonkalsrud, MD, Los Angeles

Ward O. Griffen, MD, Lexington, Ky

Donald B. Kettelkamp, MD, Indianapolis

John A. Mannick, MD, Boston

Carl H. McLauthlin, MD, Denver

Jerome H. Modell, MD, Gainesville, Fla

Samuel R. Powers, Jr., MD, Albany, NY

David C. Sabiston, Jr., MD, Durham, NC

Norman E. Shumway, Jr., MD, Stanford, Calif

Gardner W. Smith, MD, Baltimore

Richard N. Stauffer, MD, Rochester, Minn

John S. Welch, MD, Rochester, Minn

Robert Zeppa, MD, Miami, Fla

CONSULTING EDITORS

Herbert L. Abrams, MD, Boston

Viking Olov Bjork, MD, Stockholm

Robert E. Condon, MD, Milwaukee

John T. Grayhack, MD, Chicago

John B. Kinmonth, MD, London

William A. Meissner, MD, Boston

W. Keasley Welch, MD, Boston

CONSULTING EDITORS FOR AFFILIATED ORGANIZATIONS

North American Chapter of the International Cardiovascular Society:

John A. Mannick, MD, Boston

Western Surgical Association:

Robert L. Schmitz, MD, Chicago

James H. Sammons, MD
Executive Vice-President

Thomas F. Hannon
Group Vice-President,
Publishing Operations

Norman D. Richey
Director, Publication
Production Division

Helen R. Mazur
Acting Fulfillment Manager

John T. Baker, General Sales Manager;
Robert Corcoran, Eastern Sales Manager;
Fred Bauman, Richard M. Johnson, Eastern Sales Representatives; **William J. Healy**, Midwest Sales Representative; **Thomas J. Carroll**, West Coast/Northern Sales Representative; **Raymond J. Christian**, Advertising Services Manager

PUBLICATION STAFF

Offices: 535 N Dearborn St
Chicago, IL 60610

William R. Barclay, MD

Director
Scientific Publications Division

Robert W. Mayo

Executive Editor

Norman D. Richey

Editorial Processing Manager

Marlene M. Hirsch

Production Manager

Assistant Editorial Processing Manager: Diana A. Morgan

Copy Editing: Margaret Greene, Katie Sawyer, Kathi Sikora

Proofreading: Lorraine L. Kruse (Supervisor), Helen Grzeskiewicz, Madonna Niles, Elizabeth Petrikenas, Dorothy Troy, Dorothy Waier

Production Supervisors: Julia Bero (Layout), Thomas J. Handrigan (Illustrations), Marlene M. Hirsch (Editorial), James P. Bagniewski (Advertising)

Art and Senior Production Assistants: Betty Frigerio, Lucy Keating, Nancy Keyes, Shirley Latronica, JoAnne Weiskopf

Production Assistants: E. Linda Eisenberg, Anna Goins, Lucille Kuznicki, Susan M. Routhier

Manuscript Records: Mary Paino

Indexing: Phyllis H. McLaren (Supervisor), Jane P. Mauldin, Helen F. Topel

Advertising Offices.—535 N Dearborn St, Chicago, IL 60610. Area Code 312, 751-6000; and 122 E 42nd St, New York, NY 10017, Area Code 212, 867-6640.

Advertising Principles.—Each advertisement in this issue has been reviewed and complies with the principles governing advertising in AMA scientific publications. A copy of these principles is available upon request.

Subscription Rates.—The yearly rate of the ARCHIVES OF SURGERY is as follows: for members of the AMA, \$1.50, included in the annual membership dues; for non-members, \$18 for one year, \$28 for two years in the United States and US possessions; all other countries, \$28 for one year, \$38 for two years. Special yearly price to residents, interns, and medical students in the United States and US possessions, \$9.

Change of Address.—Please notify the publisher at least six weeks in advance, include both old and new address, and a mailing label taken from the most recent copy. Include your new zip code number.

THE ARCHIVES OF SURGERY is published monthly by the American Medical Association, 535 N Dearborn St, Chicago, IL 60610, and is an official publication of the Association. Second-class postage paid at Chicago, and at additional mailing office.

Please address communications concerning accepted or published communications to: AMA Specialty Journals, 535 N Dearborn St, Chicago, IL 60610.

Contents continued from p 1171.

- Chronic Testing of a Pacemaker That Needs Recharging Only Once Every Four Years** 1231
G. Frank O. Tyers, MD, FRCS(C); Robert R. Brownlee, MSEE; Howard C. Hughes, Jr, VMD; Norman J. Manley, MS; Ida N. Gorman, MS; John A. Waldhausen, MD, Hershey, Pa
Rechargeable Ag-Hg-Zn will keep up the pace.
- The Value of Scintiangiography in Arterial Disease** 1235
Charles M. Moss, MD; Amiel Z. Rudavsky, MD; Frank J. Veith, MD, New York
Has replaced contrast arteriography for screening and follow-up examination.
- Predictability of Surgical Response in Renovascular Hypertension** 1243
Joseph A. Buda, MD; Leslie Baer, MD; Jose Z. Parra-Carrillo, MD; Merhdad M. Kashef, MD; Ferdinand F. McAllister, MD; Arthur B. Voorhees, Jr, MD; Conrad L. Pirani, MD, New York
Angiotensin II blockade useful prognosticator.
- Is 80 Years Too Old For Aneurysmectomy?** 1250
Thomas F. O'Donnell, Jr, MD; R. Clement Darling, MD; Robert R. Linton, MD, Boston
Survival in unruptured group 95%.
- Abdominal Aortic Resection in Thoracic Dissection** 1258
James A. Hunter, MD; William S. Dye, MD; Hushang Javid, MD; Hassan Najafi, MD; Marshall D. Goldin, MD; Cyrus Serry, MD, Chicago
Resecting distal aorta and creating re-entry is excellent palliation for certain patients.
- "Blue Toe" Syndrome: An Indication for Limb Salvage Surgery** 1263
Allastair M. Karmody, MB, ChM, FRCS; Samuel R. Powers, MD; Vincent J. Monaco, MD; Robert P. Leather, MD, Albany, NY
Proximal lesion identified as probable source in emboli in 5 cases.
- Pediatric Vascular Trauma: Manifestations, Management, and Sequelae of Extremity Arterial Injury in Patients Undergoing Surgical Treatment** 1269
Walter M. Whitehouse, Jr, MD; Arnold G. Coran, MD; James C. Stanley, MD; Lawrence R. Kuhns, MD; William H. Weintraub, MD; William J. Fry, MD, Ann Arbor, Mich
Many injuries iatrogenic: interesting implications relative to growth if untreated.
- Arm Veins for Peripheral Arterial Reconstruction** 1276
Karl R. Clayson, MD; William H. Edwards, MD; Terry R. Allen, MD; W. Andrew Dale, MD, Nashville, Tenn
Sharper focus on indications for use of arm veins.

Contents continued on p 1174.

Emergency Carotid Artery Surgery in Neurologically Unstable Patients	1284
Jerry Goldstone, MD, Wesley S. Moore, MD, San Francisco Essential that unstable neurological deficits be managed as emergencies.	
Spontaneous Dissection of the Internal Carotid Artery	1294
William K. Ehrenfeld, MD, Edwin J. Wylie, MD, San Francisco Etiology not atherosclerosis: some may be treated conservatively.	
Simultaneous Bilateral Carotid Endarterectomies	1304
Roy H. Clauss, MD; Prafull V. Bole, MD; Mario Paredes, MD; William Doscher, MD; Adebayo Adeyemo, MD; Martin W. Kremnitzer, MD, New York Twelve patients—no neurological complications.	

REGULAR DEPARTMENTS

Books

Plastic and Reconstructive Surgery of the Breast	1311
Frank McDowell, MD, Honolulu	
Procedures in Vascular Surgery	1311
F. William Blaisdell, MD, San Francisco	
Heparin: Structure, Function and Clinical Implications	1311
Jack Lindon, PhD, Boston	
Burns of the Upper Extremity	1312
Joel M. Noe, MD, Boston	
Books Received But Not Reviewed	1312
50 Years Ago	1183
Instructions for Authors	See September 1976, p 938
Index to Advertisers	1312

Before prescribing, please consult complete product information, a summary of which follows:

INDICATIONS: Tigan is indicated for the control of nausea and vomiting.

CONTRAINDICATIONS: The injectable form of Tigan in children, the suppositories in premature or newborn infants, and use in patients with known hypersensitivity to trimethobenzamide are contraindicated. Since the suppositories contain benzocaine they should not be used in patients known to be sensitive to this or similar local anesthetics.

WARNINGS: Tigan may produce drowsiness. Patients should not operate motor vehicles or other dangerous machinery until their individual responses have been determined. *Reye's Syndrome* has been associated with the use of Tigan and other drugs, including antiemetics, although their contribution, if any, to the cause and course of the disease hasn't been established. This syndrome is characterized by an abrupt onset shortly following a nonspecific febrile illness, with persistent, severe vomiting, lethargy, irrational behavior, progressive encephalopathy leading to coma, convulsions and death.

Usage in Pregnancy: Trimethobenzamide hydrochloride was studied in reproduction experiments in rats and rabbits and no teratogenicity was suggested. The only effects observed were an increased percentage of embryonic resorptions or stillborn pups in rats administered 20 mg and 100 mg/kg and increased resorptions in rabbits receiving 100 mg/kg. In each study these adverse effects were attributed to one or two dams. The relevance to humans is not known. Since there is no adequate experience in pregnant or lactating women who have received this drug, safety in pregnancy or in nursing mothers has not been established.

PRECAUTIONS: During the course of acute febrile illness, encephalitides, gastroenteritis, dehydration and electrolyte imbalance, especially in children and the elderly or debilitated, CNS reactions such as opisthotonos, convulsions, coma and extrapyramidal symptoms have been reported with and without use of Tigan or other antiemetic agents. In such disorders caution should be exercised in administering Tigan, particularly to patients who have recently received other CNS-acting agents (phenothiazines, barbiturates, belladonna derivatives). It is recommended that severe emesis should not be treated with an antiemetic drug alone; where possible the cause of vomiting should be established. Primary emphasis should be directed toward the restoration of body fluids and electrolyte balance, the relief of fever and relief of the causative disease process. Overhydration should be avoided since it may result in cerebral edema.

The antiemetic effects of Tigan may render diagnosis more difficult in such conditions as appendicitis and obscure signs of toxicity due to overdosage of other drugs.

ADVERSE REACTIONS: There have been reports of hypersensitivity reactions and Parkinson-like symptoms. There have been instances of hypotension reported following parenteral administration to surgical patients. There have been reports of blood dyscrasias, blurring of vision, coma, convulsions, depression of mood, diarrhea, disorientation, dizziness, drowsiness, headache, jaundice, muscle cramps and opisthotonos. If these occur, the administration of the drug should be discontinued. Allergic-type skin reactions have been observed; therefore, the drug should be discontinued at the first sign of sensitization. While these symptoms will usually disappear spontaneously, symptomatic treatment may be indicated in some cases.

NOTE: The injectable form is intended for intramuscular use only; it is not recommended for intravenous use.

HOW SUPPLIED: Suppositories, Pediatric, 100 mg, boxes of 10. Suppositories, 200 mg, boxes of 10 and 50.

Ampuls, 2 ml, boxes of 10. Multiple-Dose Vials, 20 ml, Thera-Ject® (Disposable Syringes), 2 ml, boxes of 10. Capsules 250 mg, bottles of 100 and 500; 100 mg, bottles of 100.