

Copyright © 1979 by the AMERICAN MEDICAL ASSOCIATION

Official Publication, International Cardiovascular Society and Western Surgical Association

EDITORIAL

- Writing a Good Abstract Is Not Abstract Writing** 11
Arthur E. Baue, MD, New Haven, Conn

ORIGINAL ARTICLES

- Ultrasound Demonstration of Wandering Spleen** 13
Timothy G. Lee, MD; Fred E. Brickman, MD; George R. Satterwhite, MD; Leonard S. AVECILLA, MS, Portland, Ore
Ultrasound evaluation is particularly suited for patients with obscure abdominal masses.
- Anatomy and Osteotomy of the Greater Trochanter** 19
John P. Fulkerson, MD; Edmund S. Crelin, PhD; Kristaps J. Keggi, MD, Waterbury, Conn
A trochanteric osteotomy from the anterior approach during total hip replacement is described.
Editorial comment by Richard N. Stauffer, MD, Rochester, Minn
- Surgical Treatment of Adenocarcinoma:
Location: Junction of the Right, Left, and Common Hepatic Biliary Ducts** 22
Onye E. Akwari, MD, Keith A. Kelly, MD, Rochester, Minn
Resection of these tumors in selected patients prolongs survival.
- Netilmicin: A New Aminoglycoside in the Treatment of Septic Burn Patients** 28
Andrew M. Munster, MD, Baltimore
An effective new agent for organisms resistant to other aminoglycosides.
- Malignant Melanoma: Delayed Hypersensitivity Skin Testing** 35
Daniel F. Roses, MD; John F. Campion, MD; Matthew N. Harris, MD; Stephen L. Gumpert, MD, New York
Level of invasion and recurrence of disease did not correlate with skin test reactivity.
- Treatment Outcome in Recurrent Head and Neck Cancer** 39
Nathan W. Pearlman, MD, Denver
Patients identified who benefited most from further treatment.
- Gastroesophageal Fundoplication for Reflux Following Repair of
Esophageal Atresia: Experience With Nine Patients** 48
Eric W. Fonkalsrud, MD, Los Angeles
This procedure may be necessary in many children after successful atresia repair.
- Preoperative Carcinoembryonic Antigen Levels in Colorectal Carcinoma** 52
Gary Slater, MD; Angelos E. Papatestas, MD; Arthur H. Aufses, Jr, MD, New York
Preoperative levels correlated with stage of disease and tumor location.
Editorial comment by Edward Douglas Holyoke, MD, Buffalo
- Surgical Diagnosis of Mediastinal Lymphoma of Childhood** 54
Jack S. Elder, MD, Robert J. Touloukian, MD, New Haven, Conn
Lymphoma was the most frequent mediastinal mass after the age of two.
- Prosthetic Aortic Valves: Indications for and Results of Reoperation** 63
Richard J. Shemin, MD; Vincent A. Guadiani, MD; David M. Conkle, MD; Andrew G. Morrow, MD, Bethesda, Md
Reoperation performed with reasonable operative risk,
long term survival and excellent clinical improvement.

THE ARCHIVES OF SURGERY (ISSN 0004-0010) is published monthly by the American Medical Association, 535 N Dearborn St, Chicago, IL 60610, and is an official publication of the Association. Second-class postage paid at Chicago, IL 60610 and at additional mailing office.

CHANGE OF ADDRESS: POSTMASTER, send address changes to Helen R. Mazur, Director, Circulation-Fulfillment. Notification of address change should be sent at least six weeks in advance, including both old and new address and a mailing label taken from the most recent copy. Include your new zip code number.

SUBSCRIPTION RATES: Effective January 1979 the rates for the ARCHIVES OF SURGERY will be as follows: for members of the AMA, \$1.50, included in the annual membership dues; for nonmembers,

\$24 for one year, \$40 for two years in the United States and US possessions; all other countries, \$34 for one year, \$60 for two years. Special yearly price to residents, interns, and medical students in the United States and US possessions, one year, \$12; two years, \$20. Address all subscription communications to Circulation-Fulfillment Department.

ADVERTISING OFFICES: 535 N Dearborn St, Chicago, IL 60610, Area Code 312, 751-6000; and 600 Third Ave, New York, NY 10016, Area Code 212, 867-6640.

ADVERTISING PRINCIPLES: Each advertisement in this issue has been reviewed and complies with the principles governing advertising in AMA scientific publications. A copy of these principles is available on request.

CONTENTS—Continued

CLINICAL NOTES

- Postmortem Findings in Three Patients Treated With Intravenous Fat Emulsions** 66
Ib Hessov, MD; Flemming Melsen, MD; Agner Haug, MD, Århus, Denmark
Fat deposits found in the heart, and in small vessels in other organs.
Editorial comment by Basil A. Pruitt, Jr, MD, FACS, San Antonio, Tex
- Cimetidine in the Management of Metabolic Alkalosis Induced by Nasogastric Drainage** 70
Cyril H. Barton, MD; N. Dabir Vaziri, MD; Russell L. Ness, MD;
James K. Saiki, MD; Khosrow S. Mirahmadi, MD, Orange, Calif
A helpful adjunct in the treatment of severe metabolic alkalosis with renal failure.
Editorial comment by Robert Zeppa, MD, Miami
- Liver Allograft: Its Use in Chronic Active Hepatitis With Macronodular Cirrhosis, Hepatitis B Surface Antigen** 75
Jacques L. Corman, MD; Charles W. Putnam, MD; Shunzaburo Iwatsuki, MD, Denver;
Allan G. Redeker, MD, Downey, Calif; K. A. Porter, MD, London;
Robert L. Peters, MD, Los Angeles; Gerhard Schröter, MD; Thomas E. Starzl, MD, PhD, Denver
Extrahepatic infectious material was sufficient to infect the new organ.
- Salivary Gland Heterotopia in the Lower Part of the Neck** 79
William P. Adams, MD, Patricia K. Donahoe, MD, Boston
A sinus tract in an infant originated from salivary gland tissue.
- Stewart-Treves Syndrome: A Lethal Complication of Postmastectomy Lymphedema and Regional Immune Deficiency** 82
Helmut Schreiber, MD; Frank M. Barry, MD, Cleveland; W. C. Russell, MD, Middleburg Heights, Ohio;
William L. Macon IV, MD; Jeffrey L. Ponsky, MD; Walter J. Pories, MD, Cleveland
An hypothesis about etiology is proposed.
Editorial comment by Gardner W. Smith, MD, Baltimore
- Obstructing Phyto bezoar of an Intrahepatic Bile Duct** 86
Ronald A. Malt, MD, Boston
Drainage of a choledochal cyst by a cystogastrostomy resulted in formation of a bezoar.
- The Nonoperative Approach to Esophageal Perforation Due to Celestin Tube Placement** 90
Robert L. Fulton, MD; R. Neal Garrison, MD; Hiram C. Polk, Jr, MD, Louisville, Ky
The tube was used as a conduit past the site of perforation.

SPECIAL ARTICLE

- Phantom Breast Syndrome** 93
Kay Jamison, PhD; David K. Wellisch, PhD; Ronald L. Katz, MD; Robert O. Pasnau, MD, Los Angeles
Women who perceive strong emotional support from their surgeon have fewer phantom breast problems.

EXPERIMENTAL SURGERY

- Insulin and Glucagon Responses of Transplanted Intrasplenic Pancreatic Islets** 96
Bruce Nathaniel Gray; Gloria Caruso; Frank Alford; Donald Chisholm, Victoria, Australia
Greater insulin response was obtained in transplant when mechanical dissociation was used in preparation of islets.
Editorial comment by John A. Mannick, MD, Boston

TECHNIQUES

- Hand-Held Toy Squeaker During Carotid Endarterectomy in the Awake Patient** 103
Lawrence Spielberger, MD; Herman Turndorf, MD; Alfred Culliford, MD; Anthony Imparato, MD, New York
A helpful technique for neurologic monitoring.
- Removal of a Distal Common Bile Duct Stone Through Percutaneous Transhepatic Catheterization** 107
Manuel R. Perez, MD; Juan A. Oleaga, MD; David B. Freiman, MD;
Gordon L. McLean, MD; Ernest J. Ring, MD, Philadelphia
This approach may be useful when no other access is available.
Editorial comment by Hastings K. Wright, MD, New Haven, Conn

REGULAR DEPARTMENTS

- Consultants** 112
- Index to Advertisers** 111
- Instructions for Authors and Correspondents** See Dec 1978, p 1391