

PAPERS READ BEFORE THE 29TH SCIENTIFIC MEETING OF THE INTERNATIONAL CARDIOVASCULAR SOCIETY, DALLAS, JUNE 11-13, 1981—PART II

- Blood Potassium Cardioplegia Administration: Comparison of Myocardial Protection Offered by Three Techniques** 1509
 Frank P. Catinella, MD; Joseph N. Cunningham, Jr, MD;
 Ramesh K. Srungaram, MD; Ira M. Nathan, PhD;
 Edmond A. Knopp, MS; Gaetano Paone, MD; F. Gregory Baumann, PhD;
 Peter X. Adams, MD; Frank C. Spencer, MD, New York
 Multidose potassium chloride reinjection with uninterrupted aortic cross-clamping appears to be the simplest method of myocardial protection during this procedure.
- Prevention of Renal Insufficiency After Abdominal Aortic Aneurysm Resection by Optimal Volume Loading** 1517
 Harry L. Bush, Jr, MD; Jack B. Huse, MD; Willard C. Johnson, MD;
 Edward T. O'Hara, MD; Donald C. Nabseth, MD, Boston
 Pulmonary artery wedge pressure was a more accurate monitor for volume expansion than central venous pressure.
- Femorotibial and Femoroperoneal Bypass Vein Grafts: A 15-Year Experience** 1529
 George P. Kacoyanis, MD; Anthony D. Whittemore, MD;
 Nathan P. Couch, MD; John A. Mannick, MD, Boston
 Intrapopliteal arterial bypasses are associated with a 60% limb salvage rate in advanced peripheral occlusive disease.
- Sequential Femoropopliteal and Femorotibial Bypasses: A Ten-Year Follow-up Study** 1538
 Joel C. Rosenfeld, MD, Ronald P. Savarese, MD, Philadelphia; Paul Friedmann, MD, Springfield, Mass;
 Dominic A. DeLaurentis, MD, Philadelphia
 Sequential bypass grafting is a satisfactory method of revascularizing extremities when a suitable saphenous vein is not available.
- Endarterectomy for Segmental Occlusive Disease of the Superficial Femoral Artery** 1547
 Toshio Inahara, MD, Charles M. Scott, MD, Portland, Ore
 Long-term patency rates following endarterectomy of short superficial femoral artery occlusions are comparable to those following saphenous vein bypass grafts.

THE ARCHIVES OF SURGERY (ISSN 0004-0010) is published monthly by the American Medical Association, 535 N Dearborn St, Chicago, IL 60610, and is an official publication of the Association. Second-class postage paid at Chicago, IL 60610 and at additional mailing office.

SUBSCRIPTION RATES—The subscription rates for the Archives of Surgery are as follows: for members of the AMA, \$1.50 included in the annual membership dues; for nonmembers, \$30 for one year, \$56 for two years in the United States and US possessions; all other countries, one year, \$40; two years, \$76 (add \$6 surcharge to each one-year subscription, \$9 for two years for air delivery to the United Kingdom and Europe only). Special yearly rates to residents and medical students in the United States and US possessions, one year, \$15; two years, \$28. Address all subscription

communication to American Medical Association, Circulation and Fulfillment Division, 535 N Dearborn St, Chicago, IL 60610. Phone: (312)751-6079.

CHANGE OF ADDRESS—POSTMASTER, send address changes to Caryl L. Wertheimer, Director of Fulfillment, 535 N Dearborn St, Chicago, IL 60610. Notification of address change must be made at least six weeks in advance, including both old and new addresses, and a mailing label taken from the most recent copy. Include your new zip code number.

ADVERTISING PRINCIPLES: Each advertisement in this issue has been reviewed and complies with the principles governing advertising in AMA scientific publications. A copy of these principles is available on request.

CONTENTS—Continued

**Late Repair of Coarctation of the Descending Thoracic Aorta in 190 Patients:
Results up to 30 Years After Operation** 1557
Gerald M. Lawrie, MD; Michael E. DeBakey, MD; George C. Morris, Jr, MD;
E. Stanley Crawford, MD; William F. Wagner, MD; Donald H. Glaeser, DSc, Houston
The prognosis for patients undergoing repair of coarctation at the descending thoracic
aorta before age 21 years is excellent.

Postoperative Stroke and Late Neurologic Complications After Carotid Endarterectomy 1561
C. Douglas Lees, BSc, MB, ChB, Norman R. Hertzler, MD, Cleveland
Strokes involving the ipsilateral hemisphere have occurred in only 7% of patients after
carotid endarterectomy, in an average of 8.6 years of follow-up.

**Carotid Endarterectomy: The Unreliability of Intraoperative Monitoring in Patients
Having Had Stroke or Reversible Ischemic Neurologic Deficit** 1569
David Rosenthal, MD; Paul E. Stanton, Jr, MD; Pano A. Lamis, MD, Atlanta
EEG and stump pressure were unreliable indicators of cerebral blood flow in poststroke/RIND patients.

Saphenous Vein Interposition Grafts in the Microsurgical Treatment of Cerebral Ischemia 1578
Duke S. Samson, MD; Bruce L. Gewertz, MD;
Chester W. Beyer, Jr, MD; Richard M. Hodosh, MD, Dallas
The long-term patency rate of vein extracranial-intracranial bypass grafts was 80%.

Prognosis of Patients With Transient Ischemic Attacks and Normal Angiograms 1587
David S. Mendelowitz, MD; Shirley Kimmins, RN;
William E. Evans, MD, Columbus, Ohio
The prognosis for patients with normal angiograms following TIAs is good.

Obliterative Arterial Disease of the Upper Extremity 1593
Richard E. Welling, MD; John J. Cranley, MD; Raymond J. Krause, MD;
Charles D. Hafner, MD, Cincinnati
An early, aggressive approach to upper-extremity ischemic symptoms is advocated.

REGULAR DEPARTMENTS

Correspondence and Brief Communications 1597
Conservative Operative Management of Intussusception of the Appendix Norman B. Ackerman, MD, PhD, Kalkunte Suresh,
MD, Syracuse, NY . . . **Colectomy With Rectal Mucosectomy and Ileoanal Anastomosis** Max A. Heeb, MD, Sikeston, Mo

Calendar of Events 1598

Books 1598
Pernkopf Atlas of Topographical and Applied Human Anatomy (Ferner, ed)
Reviewed by Mark A. Hayes, MD, New Haven, Conn

Index to Volume 116 1599

Index to Advertisers 1592

Instructions for Authors See November 1981 issue, p 1363