

Copyright © 1981 by the AMERICAN MEDICAL ASSOCIATION

Official Publication, International Cardiovascular Society and Western Surgical Association

PAPERS READ BEFORE THE EIGHTY-EIGHTH ANNUAL MEETING OF THE WESTERN SURGICAL ASSOCIATION, SALT LAKE CITY, NOV 17-19, 1980

- The Decade Ahead: Presidential Address** 505
William H. ReMine, Rochester, Minn
- Islet Cell Autotransplantation After Pancreatectomy for Chronic Pancreatitis: Its Limitations** 511
Carlos Grodzinsky, MD; Steven Malcom, MScB; José Goldman, MD; Stanley Dienst, MD; H. K. Oh, MD; Patricia Westrick, Detroit
A major limitation may be the low yield of islet cells from the fibrotic pancreas.
- Intractable Ascites Treated With Peritoneovenous Shunts (LeVeen): A 24- to 64-Month Follow-up of Results in 52 Alcoholic Cirrhotics** 518
Herbert B. Greenlee, MD; Malcolm M. Stanley, MD; George F. Reinhardt, MD; Hines, III
Although there are many problems, the operation allows at least partial rehabilitation and ambulatory care.
- Synchronous Anterior Celiotomy and Posterior Drainage of Pancreatic Abscess** 527
Thomas V. Berne, MD, Arthur J. Donovan, MD, Los Angeles
This approach allows both adequate exploration of the abdomen and dependent drainage.
- Elective Cholecystectomy: The Role of Biliary Bacteriology and Administration of Antibiotics** 537
Michael B. Farnell, MD; Jonathan A. van Heerden, MB, FRCS(C); Robert W. Beart, Jr, MD, Rochester, Minn
Although bacteria were found in the bile of one fourth of the patients, this did not justify the use of antibiotics.
- Hypercalcemia in Patients With Breast Cancer: Osseous Metastases, Hyperplastic Parathyroid Tissue, or Pseudohyperparathyroidism?** 545
Robert C. Hickey, MD; Naguib A. Samaan, MD, PhD; Gilchrist L. Jackson, MD, Houston
Primary hyperparathyroidism may occur in patients with breast cancer and mimic osseous metastases.
- Limited Subtotal Thyroidectomy for Graves' Disease** 561
Richard D. Liechty, MD; Steven G. Silverberg, MD; James Hernandez, MD, Denver
This procedure was associated with a low incidence of recurrent hyperthyroidism.
- The Morbidity of Total Thyroidectomy** 568
Edward F. Scanlon, MD; Jacqueline E. Kellogg; David P. Winchester, MD; Richard H. Larson, MD, Evanston, Ill
Long-term follow-up of patients after total thyroidectomy indicates few problems.
- Nesidioblastosis in Adults: A Surgical Dilemma** 575
Jay K. Harness, MD, Ann Arbor, Mich; Glenn W. Geelhoed, MD, Washington, DC; Norman W. Thompson, MD, Ann Arbor, Mich;
Ronald H. Nishiyama, MD, Portland, Me; Stefan S. Fajans, MD, Richard O. Kraft, MD, Ann Arbor, Mich;
Donald R. Howard, MD, Portland, Me; Kenneth A. Clark, MD, Ann Arbor, Mich
An increase in size and number of islets throughout the pancreas was found in six adults with hyperinsulinemia.

THE ARCHIVES OF SURGERY (ISSN 0004-0010) is published monthly by the American Medical Association, 535 N Dearborn St, Chicago, IL 60610, and is an official publication of the Association. Second-class postage paid at Chicago, IL 60610 and at additional mailing office.

CHANGE OF ADDRESS: POSTMASTER, send address changes to Helen R. Mazur, Director, Circulation-Fulfillment. Notification of address change should be sent at least six weeks in advance, including both old and new address and a mailing label taken from the most recent copy. Include your new zip code number.

SUBSCRIPTION RATES: The rates for the ARCHIVES OF SURGERY are as

follows: for members of the AMA, \$1.50, included in the annual membership dues; for nonmembers, \$30 for one year, \$56 for two years in the United States and US possessions; all other countries, \$40 for one year, \$76 for two years. Special yearly price to residents, interns, and medical students in the United States and US possessions, one year, \$15; two years, \$28. Address all subscription communications to Circulation Fulfillment Department. Phone: 312, 751-6079.

ADVERTISING PRINCIPLES: Each advertisement in this issue has been reviewed and complies with the principles governing advertising in AMA scientific publications. A copy of these principles is available on request.

CONTENTS—Continued

Twenty-four-Hour pH Monitoring of Esophageal Function: Its Use in Evaluation in Symptomatic Patients After Truncal Vagotomy and Gastric Resection or Drainage	581
Gerald C. O'Sullivan, MD, MSc, FRCS(I); Tom R. DeMeester, MD; Robin B. Smith, ChM, FRCS; James W. Ryan, MD, Chicago; Lawrence F. Johnson, MD, Washington, DC; David B. Skinner, MD, Chicago	
These patients had less upright reflux and more in the supine position than patients with reflux as a primary problem.	
Simple Closure of Perforated Peptic Ulcer: Still an Effective Procedure for Patients With Delay in Treatment	591
Kenneth R. Sirinek, MD, PhD; Barry A. Levine, MD; Wayne H. Schwesinger, MD; J. Bradley Aust, MD, PhD, San Antonio, Tex	
Advanced age and delay in treatment of 30 hours or more increased morbidity and mortality.	
Cimetidine vs Antacid in Prophylaxis for Stress Ulceration	597
John A. Weigelt, MD; Christina M. Aurbakken, RN; Bruce L. Gewertz, MD; William H. Snyder III, MD, Dallas	
Antacid is more effective, safer, and less expensive.	
Gastric (Reservoir) Reduction for Morbid Obesity	602
Lawrence H. Wilkinson, MD, Ole A. Peloso, MD, Albuquerque, NM	
The stomach is wrapped with a sheet of plastic mesh to reduce its reservoir capacity.	
Endoscopic Retrograde Cholangiopancreatography: Its Use in the Evaluation of Nonjaundiced Patients With the Postcholecystectomy Syndrome	606
Marc Cooperman, MD; John J. Ferrara, MD; Larry C. Carey, MD; F. B. Thomas, MD;	
Edward W. Martin, Jr, MD; John J. Fromkes, MD, Columbus, Ohio	
This was the most useful technique for evaluating patients with symptoms after cholecystectomy.	
Factors Affecting Revision Rate After Continent Ileostomy	610
Roger R. Dozois, MD; Keith A. Kelly, MD; Duane Ilstrup, MS; Robert W. Beart, Jr, MD; Oliver H. Beahrs, MD, Rochester, Minn	
Revision was required less often in females, younger patients, and those having a continent ileostomy at the time of colectomy.	
Roux-en-Y Operation in the Management of Postoperative Fistula	614
Gerald T. Ujiki, MD, Thomas W. Shields, MD, Chicago	
A useful technique when adhesions and inflammation make resection or simple closure difficult.	
Ileocecal Valve Replacement: Its Effect on Transit Time, Survival, and Weight Change After Massive Intestinal Resection	618
Joshua Careskey, MD; Thomas R. Weber, MD; Jay L. Grosfeld, MD, Indianapolis	
Construction of a jejunal valve after massive intestinal resection in experimental animals improved survival, transit time, and weight change.	
Colectomy With Rectal Mucosectomy and Ileoanal Anastomosis in Young Patients: Its Use for Ulcerative Colitis and Familial Polyposis	623
Robert L. Telander, MD, Jean Perrault, MD, Rochester, Minn	
A feasible, although tedious, operation to provide rectal continence after colectomy.	
Hartmann's Procedure: Its Use With Complicated Carcinomas of Sigmoid Colon and Rectum	630
Stephen G. ReMine, MD, Roger R. Dozois, MD, Rochester, Minn	
A useful and safe procedure that can be associated with satisfactory long-term survival.	
Surgery, Nutritional Support, and Survival in Patients With End-Stage Renal Disease	634
Joseph L. Giacchino, MD; W. Peter Geis, MD; Bruce H. Wittenstein, MD; Vinod K. Bansal, MD;	
Vasant C. Gandhi, MD; Leonard L. Vertuno, MD, Maywood, Ill	
Nutritional support seemed helpful for patients with end-stage renal disease who required operations.	
Splenectomy for Immune Thrombocytopenic Purpura	645
Steven J. Mintz, MD; Scott R. Petersen, MD; Bruce Cheson, MD; Linda J. Cordell, MD; Ralph C. Richards, MD, Salt Lake City	
A positive response to steroids predicts a satisfactory response to splenectomy.	
Is Splenic Salvage Safe in the Traumatized Patient?	651
Armando E. Giuliano, MD, Robert C. Lim, Jr, MD, San Francisco	
Not all injured spleens can be salvaged but in selected patients such salvage is safe.	
Altered Interstitial Fluid Space Dynamics and Postresuscitation Hypertension	657
Clyde W. Dawson, MD; Charles E. Lucas, MD; Anna M. Ledgerwood, MD, Detroit	
Hypertension after resuscitation was associated with an increased plasma to interstitial fluid volume ratio.	
Cervicodiastinal Arterial Injuries: A Surgical Challenge	663
John V. Robbs, ChM, FRCS; Lynne W. Baker, MSc, FRCS; Randolph R. Human, FCS(SA);	
Ismael S. Vawda, FRCS; Harold Duncan, FRCS; P. Rajaruthnam, FRCS, Durban, South Africa	
False aneurysm was a frequent presenting sign.	

Continued on page 494.

CONTENTS—Continued

Pediatric Renal Artery Occlusive Disease and Renovascular Hypertension: Etiology, Diagnosis, and Operative Treatment 669
James C. Stanley, MD, Ann Arbor, Mich, William J. Fry, MD, Dallas
Arterial reconstructive surgery will benefit most pediatric patients with renovascular hypertension.

Immediate Jejunostomy Feeding: Its Use After Major Abdominal Trauma 681
Ernest E. Moore, MD; Ernest L. Dunn, MD; Todd N. Jones, RN, Denver
Early, safe nutritional support was provided for this group of patients.

Arterial Injuries in Children 685
J. David Richardson, MD; Mary Fallat, MD; Hirikati S. Nagaraj, MD;
Diller B. Groff, MD; Lewis M. Flint, MD, Louisville
Early exploration and repair of vascular injuries is as important in young children as it is in adults.

Selective Management of Penetrating Neck Trauma: Cost Implications 691
Robert M. Merion, MD; Jay K. Harness, MD; Stephen R. Ramsburgh, MD;
Norman W. Thompson, MD, Ann Arbor, Mich
An algorithm for selective evaluation and treatment of penetrating neck injuries is presented.

Acute and Chronic Traumatic Arteriovenous Fistulae in Civilians: Epidemiology and Treatment 697
Kenneth R. Kollmeyer, MD; John L. Hunt, MD; Brian A. Ellman, MD; William J. Fry, MD, Dallas
These lesions are frequently not clinically evident.

A University-Staffed, Private Hospital-Based Air Transport Service: The Initial Two-Year Experience 703
James H. Duke, Jr, MD, William P. Clarke, MD, Houston
The primary purpose of the program was to minimize the time to definitive treatment.

Femorotibial Bypass Grafts With Sequential Technique: Clinical Results 709
Fredric Jarrett, MD; Herbert A. Berkoff, MD; Andrew B. Crummy, MD; Folkert O. Belzer, MD, Madison, Wis
The sequential technique may increase flow and help with limb salvage.

Successful Management of Prosthetic Graft Infection With Continuous Povidone-Iodine Irrigation 716
Jack H. M. Kwaan, MD, John E. Connolly, MD, Irvine, Calif
Infarcted but functional prosthetic grafts were preserved by this method.

Popliteal Arterial Aneurysms: Their Natural History and Management 724
D. Emerick Szilagyi, MD; Richard L. Schwartz, MD; Daniel J. Reddy, MD, Detroit
Aneurysms that are 2 cm or larger in diameter should be treated by operation.

REGULAR DEPARTMENTS

Correspondence and Brief Communications 729
Femoral Incision for Arterial Bypass Adolf Singer, MD, Elmhurst, NY . . . Needle-Catheter Jejunostomy Carey P. Page, MD, San Antonio, Tex . . . Angled Low Anterior Resection Scissors E. J. Ledesma, MD; M. Tseng, MD; A. Mittelman, MD, Buffalo

Calendar of Events 732

News and Announcements 732

Index to Advertisers 566

Instructions for Authors See February 1981, p 139