

**PAPERS READ BEFORE THE FOURTH ANNUAL MEETING OF THE SURGICAL
INFECTION SOCIETY, MONTREAL, APRIL 30 TO MAY 1, 1984—PART II**

- Comparison of Indium 111-Labeled Platelets v Leukocytes In a Pyogenic Abscess** 137
Thomas P. Wright, MD; Harvey J. Sugerman, MD; James L. Tatum, MD; Jerry I. Hirsch, PharmD;
David E. Sharp, MD; G. Dastgir Qureshi, MD, Richmond, Va
Labeled leukocytes are the preferred radionuclide imaging agents for
abscess detection.
- Plasma Fibronectin Response to *Escherichia coli* and Hemoglobin** 142
Thomas S. Velky, MD; Frank Kagawa, MD; A. Gerson Greenburg, MD, PhD;
James C. Yang, MD, San Diego
Early depression of fibronectin levels occurred only when both
Escherichia coli and hemoglobin were injected in these animals.
- Lethal Microbial Synergism In Intra-abdominal Infections: *Escherichia coli* and *Bacteroides fragilis*** 146
Ori D. Rotstein, MD; Timothy L. Pruett, MD; Richard L. Simmons, MD, Minneapolis
A heat-stable surface factor produced by *Bacteroides fragilis* seems to
increase the lethal effects of *Escherichia coli*.
- Severity of Illness In Intra-abdominal Infection: A Comparison of Two Indexes** 152
Tommy Skau, MD; Per-Olof Nyström, MD, PhD; Christer Carlsson, MD, PhD, Linköping, Sweden
An acute physiology score and a septic severity score both predict risk
and could be used to evaluate treatment.
- Pulmonary Dysfunction Secondary to Soft-Tissue Endotoxin** 159
Harold Wenger, MD; Catherine Wong, MD; Robert H. Demling, MD, Boston
Endotoxin injected into soft tissues was associated with
thromboxane release, hypoxia, and pulmonary hypertension.
- The Relationship of Insulin Production to Glucose Metabolism In Severe Sepsis** 166
Michael S. Dahn, MD; Lloyd A. Jacobs, MD; Stuart Smith, PhD; Bijaya Hans, MD;
M. Patricia Lange, BSN, Allen Park, Mich; Robert A. Mitchell, PhD, John R. Kirkpatrick, MD, Detroit
Sepsis-related hyperglycemia did not appear to be associated with
peripheral insulin resistance.
- Modulation of Hepatocyte Protein Synthesis During Co-cultivation With
Macrophage-Rich Peritoneal Cells In Vitro** 180
Gary A. Keller, MD; Michael A. West, MD; John T. Harty;
Frank B. Cerra, MD; Richard L. Simmons, MD, Minneapolis
Macrophage-rich peritoneal cells, when co-cultivated with isolated hepatocytes,
significantly diminished hepatocyte protein synthesis.

THE ARCHIVES OF SURGERY (ISSN 0004-0010) is published monthly by the American Medical Association, 535 N Dearborn St, Chicago, IL 60610, and is an official publication of the Association. Second-class postage paid at Chicago, IL 60610 and at additional mailing office.

SUBSCRIPTION RATES—The subscription rates for the Archives of Surgery are as follows: for members of the AMA, \$1.50 included in the annual membership dues; for nonmembers, \$36 for one year, \$65 for two years in the United States and US possessions; all other countries, one year, \$46; two years, \$85 (add \$10 surcharge to each one-year subscription, \$20 for two years for air delivery to the United Kingdom and Europe only). Special yearly rates to residents and medical students in the United States and US possessions, one year, \$18; two years, \$32.50. Rates for subscriptions for delivery to Bangladesh, India, Japan, Nepal, South Korea, and Sri Lanka are available through respective exclusive agents. Address all subscription

communications to the American Medical Association, Circulation and Fulfillment Division, 535 N Dearborn St, Chicago, IL 60610. Phone: (312) 280-7154.

CHANGE OF ADDRESS—POSTMASTER, send all address changes to Caryl L. Wertheimer, Director of Fulfillment, 535 N Dearborn St, Chicago, IL 60610. Notification of address change must be made at least six weeks in advance, include both old and new addresses, a recent mailing label, and your new zip code.

ADVERTISING PRINCIPLES: Each advertisement in this issue has been reviewed and complies with the principles governing advertising in AMA scientific publications. A copy of these principles is available on request. The appearance of advertising in AMA publications is not an AMA guarantee or endorsement of the product or the claims made for the product by the manufacturer.

CONTENTS—Continued

Reprioritization of Hepatic Plasma Protein Release in Trauma and Sepsis	187
Gabriele Sganga, MD, Rome; John H. Siegel, MD; Glenn Brown, PhD; Bill Coleman, PhD; Charles E. Wiles III, MD; Howard Belzberg, MD; Suzanne Wedel, MD; Robert Placko, Baltimore	
Acute-phase problems increase in injured patients, with a further rise in patients in whom sepsis develops.	
In Vivo Neutrophil Delivery to Inflammatory Sites In Surgical Patients: Correlation With In Vitro Neutrophil Chemotaxis and Adherence	205
Joel S. Morris, MD; Jonathan L. Meakins, MD, DSc; Nicholas V. Christou, MD, PhD, Montreal	
Major operations lead to increased leukocyte adherence, decreased chemotaxis, and decreased delivery of leukocytes to areas of inflammation.	
Escherichia coli and Human Neutrophils: Effect of Bacterial Supernatant With Hemolysin Activity Upon Chemotaxis Receptors	212
Timothy L. Pruett, MD, Minneapolis; Dennis E. Chenoweth, MD, PhD, San Diego; Vance D. Fiegel; Robert D. Nelson, PhD; Richard L. Simmons, MD, Minneapolis	
<i>Escherichia coli</i> that produce α -hemolysin may have enhanced survival due to altering neutrophil function.	
Microbial Colonization in a New Intensive Care Burn Unit: A Prospective Cohort Study	217
Albert T. McManus, PhD; William F. McManus, MD; Arthur D. Mason, Jr, MD; Annette R. Aitchison, RN, MS; Basil A. Pruitt, Jr, MD, Fort Sam Houston, Tex	
Cohort separation may be a practical way of eliminating endemic, resistant gram-negative organisms from burn units.	
Expanded Criteria for Percutaneous Abscess Drainage	227
Stephen G. Gerzof, MD; Willard C. Johnson, MD; Alan H. Robbins, MD; Donald C. Nabseth, MD, Boston	
Percutaneous abscess drainage was effective in 82% of simple abscesses but only in 45% of complex abscesses.	
ORIGINAL ARTICLES	
Adenosine Triphosphate–Magnesium Chloride Ameliorates Reperfusion Injury Following Ischemia as Determined by Phosphorus Nuclear Magnetic Resonance	233
Bauer E. Sumpio, MD, PhD; Irshad H. Chaudry, PhD; Arthur E. Baue, MD, New Haven, Conn	
Intracellular acidosis, adenosine triphosphate (ATP) depletion, and decreased flow during reperfusion injury were rapidly reversed by ATP–magnesium chloride given after ischemia.	
Choledochoduodenostomy for Calculous Biliary Tract Disease	241
Michel Huguier, MD; François Lacaine, MD; Sidney Houry, MD; Gérard Pascal, MD, Paris	
Choledochoduodenostomy was a safe and effective operation for treatment of choledocholithiasis in older patients.	
REGULAR DEPARTMENTS	
Correspondence and Brief Communications	243
Heel-Drop Jarring Test for Appendicitis George B. Markle IV, MD, Carlsbad, NM . . . Toxic Shock Syndrome After Skin Grafting Roger E. Schmid, MD, Joseph E. Bouthillier, MD, Putnam, Conn . . . Vaginal Cuff Dehiscence Presenting as an Acute Abdomen LCDR Gregory Timberlake, MC, USN, CAPT Larry Getz, MC, USN, San Diego	
Calendar of Events	244
Books	245
Cardiac Pathology: An Integrated Text and Color Atlas (Becker, Anderson) Reviewed by Morgan Berthrong, MD, Colorado Springs, Colo . . . Vertebrobasilar Arterial Occlusive Disease: Medical and Surgical Management (Berguer, Bauer, eds) Reviewed by William Pierce, Denver . . . Anesthesia and the Cardiovascular System (Stanley, Petty, eds) Reviewed by Jessie H. Humphries, MD, Denver	
Instructions for Authors	129
Index to Advertisers	173