

Copyright 1986 by the AMERICAN MEDICAL ASSOCIATION

Official Publication of the Western Surgical Association,
Surgical Infection Society, New England Surgical Society,
Society of Surgical Oncology**PAPERS READ BEFORE THE FIFTH ANNUAL MEETING OF THE SURGICAL
INFECTION SOCIETY, NEW ORLEANS, APRIL 29 TO APRIL 30, 1985—PART II****Operative vs Percutaneous Drainage of Intra-abdominal Abscesses:****Comparison of Morbidity and Mortality 141**Jemi Olak, MD; Nicholas V. Christou, MD; Laurence A. Stein, MD;
Giovanna Casola, MD; Jonathan L. Meakins, MD, MontrealPercutaneous drainage is a reasonable initial treatment
for intra-abdominal abscesses, but other treatment may
be necessary later.**Scheduled Reoperations (Etappenlavage) for Diffuse Peritonitis 147**Wolfgang Teichmann, DH, Dietmar H. Wittmann, DH, Hamburg, West Germany;
Peter A. Andreone, MD, MinneapolisPatients with diffuse peritonitis were routinely reexplored
daily using a zipper to facilitate the operation.**Single-Antibiotic Use for Penetrating Abdominal Trauma 153**Robert M. Nelson, MD; Pamela R. Benitez, MD;
Michele A. Newell; Robert F. Wilson, MD, DetroitMoxalactam was as effective as clindamycin/tobramycin
in patients with penetrating abdominal trauma.**Trauma Serum Suppresses Superoxide Production by Normal Neutrophils 157**Marc E. Lanser, MD; Glenn E. Brown, PhD; Rafael Mora, MD;
William Coleman, PhD; John H. Siegel, MD, BaltimoreSerum from trauma patients was found to suppress the
production of superoxides and chemiluminescence in
normal white blood cells.**Antibiotics Fail to Prevent Abscess Formation Secondary to
Bacteria Trapped in Fibrin Clots 163**

Toni Hau, MD, PhD; David E. Jacobs, MD; Natalie L. Hawkins, Cleveland

Systemic antibiotics were ineffective in preventing
abscesses either because of low levels in the clots
or large numbers of organisms.

The ARCHIVES OF SURGERY (ISSN 0004-0010) is published monthly by the American Medical Association, 535 N Dearborn St, Chicago, IL 60610, and is an official publication of the Association. Second-class postage paid at Chicago, IL 60610 and at additional mailing office.

SUBSCRIPTION RATES—The subscription rates for the Archives of Surgery are as follows: for members of the AMA, \$1.50 included in the annual membership dues; for nonmembers, \$38 for one year, \$68 for two years in the United States and US possessions; all other countries, one year, \$48; two years, \$88 (add \$10 surcharge to each one-year subscription, \$20 for two years for air delivery to the United Kingdom and Europe only). Special yearly rates to residents and medical students in the United States and US possessions, one year, \$19; two years, \$34. Rates for subscriptions for delivery to Australia, Bangladesh, India, Japan, Nepal, New Zealand, South Korea, and Sri Lanka are available through respective exclusive agents.

Address all subscription communications to the American Medical Association, Circulation and Fulfillment Division, 535 N Dearborn St, Chicago, IL 60610. Phone: (312) 280-7154.

CHANGE OF ADDRESS—POSTMASTER, send all address changes to Caryl L. Wertheimer, Director of Fulfillment, 535 N Dearborn St, Chicago, IL 60610. Notification of address change must be made at least six weeks in advance, include both old and new addresses, a recent mailing label, and your new zip code.

ADVERTISING PRINCIPLES: Each advertisement in this issue has been reviewed and complies with the principles governing advertising in AMA scientific publications. A copy of these principles is available on request. The appearance of advertising in AMA publications is not an AMA guarantee or endorsement of the product or the claims made for the product by the manufacturer.

CONTENTS—Continued

Influence of Dietary Nucleotide Restriction on Bacterial Sepsis and Phagocytic Cell Function in Mice 169
Anil D. Kulkarni, MSc; William C. Fanslow; David B. Drath, PhD;
Frederick B. Rudolph, PhD; Charles T. Van Buren, MD, Houston
Nucleotides in the diet were necessary to maintain host resistance to bacterial challenge in this animal model.

Alterations in Exogenous Substrate Metabolism in Sepsis 173
Robert C. Fried, MD; Patricia M. Bailey, MD;
James L. Mullen, MD, Philadelphia; T. Peter Stein, PhD, Newark, NJ;
Lon O. Crosby, PhD, Gordon P. Buzby, MD, Philadelphia
The ability to respond appropriately to infused substrate reflected the severity of illness in the host.

Both Inflammatory and Endocrine Mediators Stimulate Host Responses to Sepsis 179
James M. Watters, MD, FRCS(C); Palmer Q. Bessey, MD;
Charles A. Dinarello, MD; Sheldon M. Wolff, MD;
Douglas W. Wilmore, MD, Boston
Injection of an inflammatory agent combined with infusion of the counter-regulatory hormones produced in normal volunteers the host responses to critical illness.

Oxygen as an Antibiotic: A Comparison of the Effects of Inspired Oxygen Concentration and Antibiotic Administration on In Vivo Bacterial Clearance 191
David R. Knighton, MD; Betty Halliday;
Thomas K. Hunt, MD, San Francisco
The combination of increased inspired oxygen and antibiotic administration was additive and eliminated almost all infectious necrosis in this model.

PANEL DISCUSSION—SURGICAL INFECTION SOCIETY 1985

Multiple-Organ-Failure Syndrome 196
C. James Carrico, MD, Seattle; Jonathan L. Meakins, MD, DSc, FRCSC, FACS,
J. C. Marshall, MD, FRCSC, Montreal; Donald Fry, MD, Cleveland; Ronald V. Maier, MD, Seattle

ORIGINAL ARTICLES

Regional Brain Amino Acid and Neurotransmitter Derangements During Abdominal Sepsis and Septic Encephalopathy in the Rat: The Effect of Amino Acid Infusions 209
Herbert R. Freund, MD; Michael Muggia-Sullam, MD; Richard LaFrance, PharmD;
Jane Holroyde; Josef E. Fischer, MD, Cincinnati
Alterations in brain neurotransmitter metabolism may be related to the development of septic encephalopathy.

Successful Vein Bypass in Patients With an Ischemic Limb and a Palpable Popliteal Pulse 217
Nancy L. Cantelmo, MD; J. Robert Snow, MD; James O. Menzoian, MD;
Frank W. LoGerfo, MD, Boston
The results of distal vein graft reconstruction from the superficial femoral or popliteal artery are comparable to results with more proximal disease.
Editorial Comment by Calvin B. Ernst, MD, Detroit

Continued on page 130.

One-Stage Simple Mastectomy With Immediate Reconstruction for High-Risk Patients: An Improved Technique: The Biologic Basis for Ductal-Glandular Mastectomy	221
Kirby I. Bland, MD; Baron O'Neal, MD; Leonard J. Weiner, MD; Gordon R. Tobin II, MD, Louisville	
This technique is an alternative to simple mastectomy in women with a high risk for developing breast cancer. Invited Editorial Comment by Stephan Ariyan, MD, New Haven, Conn	
Double-Lumen, Silicone Rubber, Indwelling Venous Catheters: A New Modality for Angioaccess	229
Harry Schanzer, MD; Steven Kaplan, MD; Juan Bosch, MD; Sheldon Glabman, MD; Lewis Burrows, MD, New York	
Major advantages of this technique include the elimination of the need for venipuncture and sacrifice of arteries.	
Necrotizing Fasciitis	233
David Gozal, MD; Avishai Ziser, MD; Avigdor Shupak, MD; Amnon Ariel, MD; Yehuda Melamed, MD, Haifa, Israel	
Sixteen patients with necrotizing fasciitis were treated by a uniform protocol of radical excision, antibiotics, and hyperbaric oxygen, with two deaths occurring.	
Jejunostomy: A Rarely Indicated Procedure	236
Mark B. Adams, MD, MS; Gary R. Seabrook, MD; Edward A. Quebbeman, MD, PhD; Robert E. Condon, MD, MS, Milwaukee	
Jejunostomy had a substantial risk of mortality and of complications.	
The Effects of Perioperative Fluorouracil Administration on Convalescence and Wound Healing	239
Joseph M. Klausner, MD; Shlomo Lelcuk, MD; Moshe Inbar, MD; Ronald Rozin, MD, Tel Aviv, Israel	
The administration of fluorouracil during and after operation had no deleterious effect on wound healing and recovery. Editorial Comment by LaSalle D. Leffall, Jr, MD, Washington, DC	
CLINICAL NOTES	
Esophageal Perforation After Fiberoptic Endoscopic Injection Sclerotherapy for Esophageal Varices	243
Elijahou Shemesh, MD, Leon Bat, MD, Tel Hashomer, Israel	
Perforation occurred in two patients (1% of all procedures) and both were treated successfully without operation.	
Chylous Ascites Following Resection of a Ruptured Abdominal Aneurysm: Treatment With a Peritoneovenous Shunt	246
William G. Sarazin, MD, Kendall E. Sauter, MD, MS, Milwaukee	
After a low-fat diet, hyperalimentation, and paracenteses were unsuccessful, a peritoneovenous shunt provided for recovery.	
REGULAR DEPARTMENTS	
Instructions for Authors	136
Index to Advertisers	248