

EDITORIAL

- Who Is Captain of the Anesthesia Ship?** 753
Jerome H. Modell, MD, Gainesville, Fla
- PAPERS READ BEFORE THE 12TH ANNUAL MEETING OF THE NEW ENGLAND SOCIETY FOR VASCULAR SURGERY, WHITEFIELD, NH, OCT 10-12, 1985**
- Can Ventilation-Perfusion Scans Accurately Diagnose Acute Pulmonary Embolism?** 754
Kevin McBride, MD; Wayne W. La Morte, MD, PhD; James O. Menzoian, MD, Boston
A high error rate would have resulted if only ventilation-perfusion scans were relied on to diagnose pulmonary embolism.
- Recognition and Management of Impending Vein-Graft Failure: Importance for Long-term Patency** 758
Jon R. Cohen, MD; John A. Mannick, MD; Nathan P. Couch, MD; Anthony D. Whittemore, MD, Boston
Significant postoperative reduction in Doppler ankle/brachial index should lead to evaluation and therapy for stenotic lesions.
- Effect of Prolonged Intra-arterial Prostaglandin E₁ Infusion on Canine Arterial Graft Patency** 760
Jack L. Cronenwett, MD, Hanover, NH; David M. Shapiro, MD, Joshua L. Luce, Ann Arbor, Mich
Local infusion of PGE₁ increased limb blood flow, but improved graft patency seemed to be due to a systemic effect.
- Hemodynamic Indicators of Postthrombotic Sequelae** 765
C. Scott Norris, MD, Jeffrey M. Darrow, MD, Worcester, Mass
Noninvasive testing after acute deep venous thrombosis may identify patients requiring elastic compression therapy.
- Intra-arterial Fibrinolytic Therapy: Efficacy of Streptokinase vs Urokinase** 769
Michael Belkin, MD; Barbara Belkin, MD; Charles A. Bucknam, MD; J. John Straub, MD; Robert Lowe, MD, Hartford, Conn
Urokinase was much more effective and safer than streptokinase but was also more expensive.
- Deep-Penile-Vein Arterialization for Arterial and Venous Impotence** 774
Alexander Balko, MD; Chander M. Malhotra, MD; John P. Wincze, PhD; Jacques G. Susset, MD;
Sudhir Bansal, MD; Wilfred I. Carney, MD; Robert W. Hopkins, MD, Providence, RI
This technique was beneficial for patients with impotence due to venous leakage from the corpora cavernosa bodies during erection.
- Antiplatelet Therapy and Vascular Grafts: Studies in a Baboon Ex Vivo Shunt** 778
Jens Eldrup-Jorgensen, MD; Raymond J. Connolly, PhD; William C. Mackey, MD; Karen Ramberg, MS;
Thomas F. O'Donnell, MD; C. Robert Valeri, MD; Allan D. Callow, MD, PhD, Boston
Both aspirin and heparin inhibited platelet deposition on grafts in this model.
- Efficacy of Tissue Plasminogen Activator and Urokinase in a Canine Model of Prosthetic Graft Thrombosis** 782
G. Richard Curl, MD; Joseph A. Jakubowski, PhD; Donald C. Nabseth, MD; Harry L. Bush, Jr, MD, Boston
Tissue plasminogen activator and urokinase produced lysis of clots, but extravasation of blood occurred only with the former.

ORIGINAL ARTICLES

- Energy Expenditure in Malnourished Patients With Colorectal Cancer** 789
Daniel T. Dempsey, MD; Linda S. Knox, BSN; James L. Mullen, MD; Clifford Miller, MS;
Irene D. Feurer, MS; Gordon P. Buzby, MD, Philadelphia, Pa
Energy expenditure did not correlate with tumor burden but may be related to duration of disease.
- Murine Neutrophil Migration During the Course of an Intra-abdominal Abscess** 797
William Lineaweaver, MD; David Soucy; Richard Howard, MD, PhD, Gainesville, Fla
Neutrophil migration increased initially and then was depressed in this intra-abdominal abscess mouse model.
Editorial Comment by David L. Dunn, MD, PhD, Richard L. Simmons, MD, Minneapolis

The ARCHIVES OF SURGERY (ISSN 0004-0010) is published monthly by the American Medical Association, 535 N Dearborn St, Chicago, IL 60610, and is an official publication of the Association. Second-class postage paid at Chicago, IL 60610 and at additional mailing office.

SUBSCRIPTION RATES—The subscription rates for the Archives of Surgery are as follows: for members of the AMA, \$1.50 included in the annual membership dues; for nonmembers, \$38 for one year, \$68 for two years in the United States and US possessions; all other countries, one year, \$48; two years, \$88 (add \$10 surcharge to each one-year subscription, \$20 for two years for air delivery to the United Kingdom and Europe only). Special yearly rates to residents and medical students in the United States and US possessions, one year, \$19; two years, \$34. Rates for subscriptions for delivery to Australia, Bangladesh, India, Japan, Nepal, New Zealand, South Korea, and Sri Lanka are available through respective exclusive agents.

Address all subscription communications to the American Medical Association, Circulation and Fulfillment Division, 535 N Dearborn St, Chicago, IL 60610. Phone: (312) 280-7154.

CHANGE OF ADDRESS—POSTMASTER, send all address changes to Caryl L. Wertheimer, Director of Fulfillment, 535 N Dearborn St, Chicago, IL 60610. Notification of address change must be made at least six weeks in advance, include both old and new addresses, a recent mailing label, and your new zip code.

ADVERTISING PRINCIPLES: Each advertisement in this issue has been reviewed and complies with the principles governing advertising in AMA scientific publications. A copy of these principles is available on request. The appearance of advertising in AMA publications is not an AMA guarantee or endorsement of the product or the claims made for the product by the manufacturer.

CONTENTS—Continued

Abdominal Incisions and Staging in Ovarian Cancer	800
Larry McGowan, MD, Washington, DC	
Incomplete intraoperative evaluation of the extent of disease was found in many patients with ovarian cancer who had transverse incisions.	
Invited Editorial Comment by John McL. Morris, MD, New Haven, Conn	
Aseptic Hip Necrosis After Renal Transplantation	803
Maria Brazil, MD; Robin J. Linderer, RN; Maureen J. Dickhans, RN; Paul J. Garvin, MD, St Louis	
Renal allograft dysfunction may contribute to the development of aseptic hip necrosis.	
Mammography of Women With Suspicious Breast Lumps	807
Jeremiah O. Young, MD, Framingham, Mass; Norman L. Sadowsky, MD, Jamaica Plain, Mass;	
John W. Young, New Haven, Conn; Leon Herman, MD, Framingham, Mass	
Preoperative mammography is indicated to screen for occult carcinoma, to prevent biopsy sampling errors and to help estimate the adequacy of lumpectomy.	
Antacid vs Sucralfate in Preventing Acute Gastrointestinal Tract Bleeding	
In Abdominal Aortic Surgery: A Randomized Trial in 50 Patients	810
Edgar Borrero, MD; Joy Ciervo, RN; John B. Chang, MD, New Hyde Park, NY	
In 25 patients receiving sucralfate, only one had upper gastrointestinal tract bleeding.	
Presentation of Congenital Diaphragmatic Hernia Past the Neonatal Period	813
Barry M. Newman, MD, Charlottesville, Va; Ehsan Afshani, MD; Melvyn P. Karp, MD;	
Theodore C. Jewett, Jr, MD; Donald R. Cooney, MD, Buffalo	
This diagnosis should be considered in children with persistent gastrointestinal or respiratory problems and an abnormal chest x-ray film.	
Invited Editorial Comment by Robert J. Touloukian, MD, New Haven, Conn	
Peritoneoscopy in Extrahepatic Abdominal Diseases	818
Ajay K. Kriplani, MS, Lalit K. Sharma, MS, FICS, New Delhi, India	
Laparotomy was not necessary in two thirds of patients in whom peritoneoscopy was performed.	
Editorial Comment by Robert E. Hermann, MD, Cleveland	
Continuous Absorbable vs Interrupted Nonabsorbable Fascial Closure: A Prospective, Randomized Comparison	821
Paul M. McNeill, MD, Harvey J. Sugerman, MD, Richmond, Va	
In obese patients there was little difference in results if abdominal fascial closure was by continuous absorbable or interrupted nonabsorbable sutures.	
Canine Mucosal Antrectomy	824
James M. Becker, MD; Charles B. Rosen, MD; Keith A. Kelly, MD; Paul E. Micevych, MD;	
Brian Cranley, MD; Vay Liang W. Go, MD, Rochester, NY	
Mucosal antrectomy decreased serum gastrin and gastric hydrochloride output, but this was counteracted in part by residual and new G cells.	
Editorial Comment by Lloyd M. Nyhus, MD, Chicago	
Ranitidine and Somatostatin: Their Effects on Bleeding From the Upper Gastrointestinal Tract	833
Nicola Basso, MD; M. Bagarani, MD; F. Bracci, MD; G. Cucchiara, MD; D. Gizzonio, MD;	
G. Grassini, MD; M. Percoco, MD; F. Procacciante, MD; F. Toti, MD, Rome	
Ranitidine or somatostatin did not alter the course of patients with upper gastrointestinal tract bleeding.	
Long-Acting β-Adrenergic Antagonists as Preparation for Surgery in Thyrotoxicosis	838
Paul H. Gerst, MD; John Fildes, MD; Peter Baylor, MD; Joel Zonszein, MD, New York	
Propranolol was an effective, fast-acting, and economical preparation for thyroidectomy for Graves' disease.	
Editorial Comment by Robert E. Hermann, MD, Cleveland	
Parathyroid Hormone Assay: Unreliable and Overused	841
Thomas A. Broughan, MD; Mark T. Jaroch, MD; Caldwell B. Esselstyn, Jr, MD, Cleveland	
A serum calcium determination remains the best test for diagnosing primary hyperparathyroidism.	
Radionuclide Esophageal Transit: A Screening Test for Esophageal Disorders	843
David Netscher, MD; Gerald M. Larson, MD; Hiram C. Polk, Jr, MD, Louisville	
The radionuclide esophageal transit test was as sensitive as manometry in identifying motility disorders.	
CLINICAL NOTES	
Hepatocellular Carcinoma Presenting With Intractable Diarrhea: A Radiologic-Pathologic Correlation	849
Edward Steiner, MD; Paul Velt, MD; Oscar Gutierrez, MD; Seymour Schwartz, MD; William Chey, MD, Rochester, NY	
The tumor cells were found to contain vasoactive intestinal polypeptide, gastrin, and prostaglandinlike immunoactivity.	
Pulmonary Dysfunction Associated With Streptokinase Therapy	852
Morris D. Kerstein, MD, Michael F. Adinolfi, MD, New Orleans	
The adult respiratory distress syndrome developed 48 hours after beginning an intra-arterial infusion of streptokinase.	
REGULAR DEPARTMENTS	
Books	854
Correspondence and Brief Communications	859
Calendar of Events	860
Instructions for Authors	749
Index to Advertisers	750