

**PAPERS READ BEFORE THE 6TH ANNUAL MEETING OF THE SURGICAL INFECTION SOCIETY,
CHICAGO, APRIL 21-22, 1986—PART I**

- Old Problems, New and Persistent Challenges: Presidential Address** 15
J. Wesley Alexander, MD, ScD, Cincinnati
- Physiology and Metabolism in Isolated Viral Septicemia:
Further Evidence of an Organism-Independent, Host-Dependent Response** 21
Clifford Scott Deutschman, MD, Minneapolis;
Frank N. Konstantinides, St Paul, Minn; Michael Tsai, PhD;
Richard L. Simmons, MD; Frank B. Cerra, MD, Minneapolis
The systemic response of patients with viral infections
was similar to that due to bacteria or fungi.
- Complement-Mediated Neutrophil Activation in Sepsis- and Trauma-Related Adult
Respiratory Distress Syndrome: Clarification With Radioaerosol Lung Scans** 26
Steven D. Tennenberg, MD; Martin P. Jacobs, MD;
Joseph S. Solomkin, MD, Cincinnati
Pulmonary alveolar capillary membrane permeability increased
in patients who developed adult respiratory distress syndrome.
- Microbiologic Prediction of Abdominal Surgical Wound Infection** 33
Alan V. Pollock, MB, FRCS, Mary Evans, Scarborough, North Yorkshire, England
Cultures incubated immediately in broth in the operating room
most accurately predicted the likelihood of wound infection.
- Surface Biofilm Disruption:
Enhanced Recovery of Microorganisms From Vascular Prostheses** 38
David F. Tollefson, MD; Dennis F. Bandyk, MD; Hermann W. Kaebnick, MD;
Gary R. Seabrook, MD; Jonathan B. Towne, MD, Milwaukee, Wis
The formation of an adherent bacteria biofilm may be an
important factor in late graft infection.
- Biliary Bacteria, Antibiotic Use, and Wound Infection in
Surgery of the Gallbladder and Common Bile Duct** 44
Ronald T. Lewis, MB, FRCS(C); R. Graydon Godall, MD, FRCS(C);
Breen Marien, MD, FRCS(C); Myung Park, MD, FRCS(C);
Walter Lloyd-Smith, MD, FRCS(C);
Frederick M. Wiegand, MD, FRCS(C), Montreal
The efficacy of preoperative antibiotics in high-risk patients
having biliary tract operations is documented.
- Role of the Macrophage in the Translocation of Intestinal Bacteria** 48
Carol L. Wells, PhD; Michael A. Maddaus, MD;
Richard L. Simmons, MD, Minneapolis
Mice considered macrophage-defective had decreased translocation
of intestinal bacteria to mesenteric lymph nodes.
- Effect of Blood Transfusions on Immune Function:
III. Alterations in Macrophage Arachidonic Acid Metabolism** 56
J. Paul Waymack, MD; Lois Gallon; Uno Barcelli, MD;
Orrawin Trocki, MS, RD; J. Wesley Alexander, MD, ScD, Cincinnati
Allogenic transfusions were associated with decreased
macrophage migration and production of prostaglandin E.

The **ARCHIVES OF SURGERY** (ISSN 0004-0010) is published monthly by the American Medical Association, 535 N Dearborn St, Chicago, IL 60610, and is an official publication of the Association. Second-class postage paid at Chicago, IL 60610 and at additional mailing office.

SUBSCRIPTION RATES—The subscription rates for the Archives of Surgery are as follows: for members of the AMA, \$1.50 included in the annual membership dues; for nonmembers, \$38 for one year, \$68 for two years in the United States and US possessions; all other countries, one year, \$50; two years, \$92 (add \$15 surcharge to each one-year subscription, \$30 for two years for air delivery to the United Kingdom, Europe, Bangladesh, India, Nepal, or Sri Lanka). Special yearly rates to residents and medical students in the United States and US possessions, one year, \$19; two years, \$34. Rates for subscriptions for delivery to Australia, Japan, New Zealand, or South Korea are available through respective exclusive agents. Address all

subscription communications to the American Medical Association, Circulation and Fulfillment Division, 535 N Dearborn St, Chicago, IL 60610. Phone: (312) 280-7154.

CHANGE OF ADDRESS—POSTMASTER, send all address changes to Caryl L. Wertheimer, Director of Fulfillment, 535 N Dearborn St, Chicago, IL 60610. Notification of address change must be made at least six weeks in advance, include both old and new addresses, a recent mailing label, and your new zip code.

ADVERTISING PRINCIPLES: Each advertisement in this issue has been reviewed and complies with the principles governing advertising in AMA scientific publications. A copy of these principles is available on request. The appearance of advertising in AMA publications is not an AMA guarantee or endorsement of the product or the claims made for the product by the manufacturer.

CONTENTS—Continued

Hemorrhage Without Tissue Trauma Produces Immunosuppression and Enhances Susceptibility to Sepsis 62
Rabie N. Stephan, MD; Thomas S. Kupper, MD; Alexander S. Geha, MD;
Arthur E. Baue, MD; Irshad H. Chaudry, PhD
The mitogen response and mixed lymphocyte reaction were suppressed for five days after an arterial bleed.

C-Reactive Protein Levels Predict Postoperative Septic Complications 69
Robert A. Mustard, Jr, MD, FRCS(C); John M. A. Bohnen, MD, FRCS(C);
Sabeha Haseeb, MD; Rafal Kasina, MD, Toronto
A rise in C-reactive protein levels after operation and a fall to normal after the second day with no secondary rise indicates that sepsis will not develop.

Comparison of Quantitative Microbiology and Histopathology in Divided Burn-Wound Biopsy Specimens 74
Albert T. McManus, PhD; Seung H. Kim, MD; William F. McManus, MD;
Arthur D. Mason, Jr, MD; Basil A. Pruitt, Jr, MD, Fort Sam Houston, Tex
Quantitative microbiology is not a substitute for histologic examination in the diagnosis of invasive infection.

Supplemental Dietary Tyrosine in Sepsis and Acute Hemorrhagic Shock 78
Ronald Simon, MD; Warren Wetzal, MD; Kim Winsey, MD;
Stanley M. Levenson, MD; Achilles A. Demetriou, MD, PhD, Bronx, NY
Dietary supplementation with tyrosine enhanced tolerance to intra-abdominal sepsis and hemorrhage.

Skeletal Microcirculatory Responses to Hyperdynamic Escherichia coli Sepsis in Unanesthetized Rats 86
Henry M. Cryer, MD; R. Neal Garrison, MD; Herman W. Kaebnick, MD;
Patrick D. Harris, PhD; Lewis M. Flint, MD, Louisville, Ky
With sepsis there was dilation of small terminal arterioles in skeletal muscle.

Succinic Acid Production by Bacteroides fragilis: A Potential Bacterial Virulence Factor 93
Ori D. Rotstein, MD, Toronto; Carol L. Wells, PhD;
Timothy L. Pruett, MD; Jennifer J. Sorenson;
Richard L. Simmons, MD, Minneapolis
Bacteroides fragilis produces a soluble leukocyte inhibitory factor that could be succinic acid.

Inhibition of Early Events in the Human T-Lymphocyte Response to Mitogens and Alloantigens by Hydrogen Peroxide 99
Brian M. Freed, MA; Robert Rapoport; Neil Lempert, MD, Albany, NY
Hydrogen peroxide, which can be produced by phagocytes, is an in vitro inhibitor of T-cell responses.

Macrophages and Translymphatic Absorption Represent the First Line of Host Defense of the Peritoneal Cavity 105
David L. Dunn, MD, PhD; Roderick A. Barke, MD; David C. Ewald;
Richard L. Simmons, MD, Minneapolis
Lymphatic clearance accounted for the greatest amount of bacterial removal initially followed by macrophage phagocytosis.

Increased Susceptibility to Infection due to Infusion of Exogenous Chemotaxin 111
Roger W. Yurt, MD, G. Tom Shires, MD, New York
Animals infused with a chemotaxin had a shorter survival time after a contaminated burn wound.

REGULAR DEPARTMENTS

Letters to the Editor 118

Books 123

Calendar 123

Instructions for Authors 11

Index to Advertisers 12