

**PAPERS READ BEFORE THE ANNUAL MEETING OF THE SOCIETY OF SURGICAL ONCOLOGY,  
LONDON, APRIL 27 TO APRIL 30, 1987—PART II**

- Colorectal Cancer: Dukes' Stage, Tumor Site, Preoperative Plasma CEA Level, and Patient Prognosis Related to Tumor DNA Ploidy Pattern** ..... 1375  
Nigel A. Scott, MB; Harry S. Wieand, PhD; Charles G. Moertel, MD; Stephen S. Cha, MS;  
Robert W. Beart, MD; Michael M. Lieber, MD, Rochester, Minn  
In colorectal cancer nondiploid DNA patterns were associated with decreased survival.
- Surgical Treatment of Lymph Nodes With Metastatic Melanoma From Unknown Primary Site** ..... 1380  
Jan H. Wong, MD; Leslie A. Cagle, MD; Donald L. Morton, MD, Los Angeles  
Failure to find a primary melanoma did not decrease the survival of patients treated for nodal metastases of melanoma.
- Human Immune Response to Monoclonal Antibody Administration Is Dose-Dependent** ..... 1384  
Henry F. Sears, MD, Darius J. Bągli, MD, CM, Boston; Dorothee Herlyn, DVM; Elaine DeFreitas, PhD;  
H. Suzuki, PhD, Philadelphia; Glenn Steele, MD, PhD, Boston; Hilary Koprowski, MD, Philadelphia  
A high dose of mouse monoclonal antibody produced a type of tolerance for up to five months in some patients with tumors.
- The Current Evaluation of Nonpalpable Breast Lesions** ..... 1389  
Nicholas P. Lang, MD; Gary E. Talbert, MD; Kris B. Shewmake, MD; Wilma C. Diner, MD;  
David Weiss, MD; Frank G. Bivins; Kent C. Westbrook, MD, Little Rock, Ark  
Guidelines for which mammographic findings require biopsy are presented.
- Intraoperative Single-Dose Radiotherapy: Observations on Staging and Interstitial Treatment of Unresectable Liver Metastases** ..... 1392  
Russell J. Nauta, MD; Edward K. Heres, MD; David S. Thomas, MD; K. William Harter, MD; James E. Rodgers, PhD;  
Richard W. Holt, MD; Thomas C. Lee, MD; Daniel B. Walsh, MD; Anatoly Dritschilo, MD, Washington, DC  
Intraoperative brachytherapy for liver metastases was associated with a fall in CEA values in some patients.
- The Acute Metabolic Effects of Tumor Necrosis Factor Administration in Humans** ..... 1396  
Robert S. Warren, MD; H. Fletcher Starnes, Jr, MD; Janice L. Gabrielove, MD;  
Herbert F. Oettgen, MD; Murray F. Brennan, MD, New York  
Tumor necrosis factor (cachectin) induced acute alterations in amino acid metabolism in patients with cancer.
- Composite Pelvic Resection: An Approach to Advanced Pelvic Cancer** ..... 1401  
Harold J. Wanebo, MD; Richard Whitehill, MD; Douglas Gaker, MD; Gwo J. Wang, MD;  
Raymond Morgan, MD; William Constable, MD, Charlottesville, Va  
Resection of a portion of the musculoskeletal pelvic wall for extensive pelvic cancers helped provide good local control.
- Human T Cells Specifically Activated Against Autologous Malignant Melanoma** ..... 1407  
Craig L. Slingluff, MD; Timothy L. Darrow, PhD; H. F. Seigler, MD, Durham, NC  
Lysis of autologous melanoma cells was produced by specific melanoma- and interleukin 2-activated lymphocytes.
- RAS Oncogene p21 Levels Parallel Malignant Potential of Different Human Colonic Benign Conditions** ..... 1414  
Fabrizio Michelassi, MD; Steve Leuthner; Mark Lubienski, MD; David Bostwick, MD;  
Jennifer Rodgers, HT; Mark Handcock; George E. Block, MD, Chicago  
The level of protein product of this family of genes correlated with the malignant potential of benign human colonic disease.
- In Vitro Killing of Melanoma by Liposome-Delivered Intracellular Irradiation** ..... 1417  
Sigismund S. Pikul II, MD; Norris J. Parks, PhD; Philip D. Schneider, MD, PhD, Davis, Calif  
Radioactive particles encapsulated in liposomes killed melanoma cells in vitro.
- Major Intestinal Complications of Radiotherapy: Management and Nutrition** ..... 1421  
Mervyn Deitel, MD, Toan B. To, MD, Toronto  
Resection to normal bowel, omental wrap of anastomoses, and long-tube orientation seemed helpful.

The ARCHIVES OF SURGERY (ISSN 0004-0010) is published monthly by the American Medical Association, 535 N Dearborn St, Chicago, IL 60610, and is an official publication of the Association. Second-class postage paid at Chicago, IL 60610 and at additional mailing office.

**SUBSCRIPTION RATES**—The subscription rates for the Archives of Surgery are as follows: \$40 for one year, \$70 for two years in the United States and US possessions; all other countries, one year, \$52; two years, \$96 (add \$15 surcharge for each one-year subscription, \$30 for two years for air delivery to the United Kingdom, Europe, Bangladesh, India, Nepal, or Sri Lanka). Rates for subscriptions for delivery to Australia, Japan, New Zealand, or South Korea are available through respective agents. **Special rates for residents and medical students in the United States and US possessions are available.** Address inquiries to the American Medical

Association, Circulation and Fulfillment Division, 535 N Dearborn St, Chicago, IL 60610. Phone: (312) 280-7168.

**CHANGE OF ADDRESS**—POSTMASTER, send all address changes to Caryl L. Wertheimer, Director of Fulfillment, 535 N Dearborn St, Chicago, IL 60610. Notification of address change must be made at least six weeks in advance, include both old and new addresses, a recent mailing label, and your new zip code.

**ADVERTISING PRINCIPLES:** Each advertisement in this issue has been reviewed and complies with the principles governing advertising in AMA scientific publications. A copy of these principles is available on request. The appearance of advertising in AMA publications is not an AMA guarantee or endorsement of the product or the claims made for the product by the manufacturer.

## CONTENTS—Continued

<b>Localization of Radiolabeled Monoclonal Antibody in Colorectal Cancer: Initial Dosimetry Results</b> .....	1425
Kenneth H. Cohn, MD; Sydney Welt, MD; William P. Banner, MD; Martin Harrington, MD; Samuel Yeh, MD; Junichi Sakamoto, MD; Carlos Cardon-Cardo, MD, PhD; John Daly, MD; Nancy Kemeny, MD; Alfred Cohen, MD; Kenneth Lloyd, PhD; Jerome DeCasse, MD, PhD; Herbert Oettgen, MD; Lloyd Old, MD, New York Imaging of metastatic colorectal cancer is possible with monoclonal antibody.	
<b>Neoadjuvant Chemotherapy Before Definitive Treatment for Stage III Carcinoma of the Breast</b> .....	1430
Gordon F. Schwartz, MD; Ronald I. Cantor, MD; William A. Biermann, MD, Philadelphia Preoperative chemotherapy followed by conventional treatment was associated with considerable improvement in these patients.	
<b>Expression of Autoantigens in Human Colorectal Carcinomas: Implications for a Generic Vaccine</b> .....	1435
J. Milburn Jessup, MD; Kefeng Qi, MB; Kristine Kanellopoulos; Debra E. Campbell; Karen R. Cleary, MD; Catherine M. Hickey; Christopher L. Reading, PhD; Howard E. Savage, PhD, Houston A generic vaccine for colorectal cancer may be possible because of limited dominant antigens.	
<b>Management of Patients With Clinical Stage I or II Nonseminomatous Germ Cell Tumors of the Testis: Evolving Therapeutic Options</b> .....	1443
Jerome P. Richie, MD; Mark A. Socinsky, MD; Claire Y. Fung, MD; Gilbert L. Brodsky, MD; Leslie A. Kalish, PhD; Marc B. Garnick, MD, Boston Therapy should be modified according to histologic features of the tumor, retroperitoneal disease, and tumor markers.	
<b>Natural Killer Activity of Lymphocytes Infiltrating Human Lung Cancers Following Preoperative Systemic Recombinant Interleukin 2</b> .....	1446
Timothy M. Anderson, MD; Yukihiko Ibayashi, MD; Yutaka Tokuda, MD; Steve Colquhoun, MD; Carmack Holmes, MD; Sidney H. Golub, PhD, Los Angeles Depressed natural killer lymphocyte cytotoxicity of tumor- infiltrating lymphocytes was augmented by interleukin 2.	
<b>The Impact of Delayed Surgery on Radiotherapy Dose and Local Control of Rhabdomyosarcoma</b> .....	1451
Erlinda Etcubanas, MD; Bhaskar N. Rao, MD; Larry E. Kun, MD; Marc E. Horowitz, MD; David M. Parham, MD; H. Omar Hustu, MD; Alexander A. Green, MD, Memphis Initial chemotherapy, delayed operation, and then radiation therapy helped with local tumor control.	
<b>Immunotherapy of Pulmonary Metastases Using Monoclonal Antibody to T-Cell Suppressor Factor and Interleukin 2</b> .....	1455
Benjamin Kim, MD, Patricia Warnaka, Cleveland; Michael Iverson, PhD, La Jolla, Calif; Anthony L. Imbembo, MD, Cleveland Interleukin 2 and a monoclonal antibody that inhibits T-lymphocyte suppressor activity decreased methylcholanthrene-produced pulmonary metastases.	
<b>Positive Relationship of Clinical and Serologic Responses to Vaccinia Melanoma Oncolysate</b> .....	1460
Marc K. Wallace, MD, Jerry A. Bash, PhD, Miami Beach, Fla; Eleuthere Leftheriotis, DVM, Lyon, France; Hilliard Seigler, MD, Durham, NC; Kirby Bland, MD, Gainesville, Fla; Harold Wanebo, MD, Charlottesville, Va; Charles Baich, MD, Alfred A. Bartolucci, PhD, Birmingham, Ala In patients with stage II melanoma, vaccinia melanoma oncolysate increased disease-free survival when compared with historical controls.	
<b>Osteosarcoma and Other Neoplasms of Bone: Magnetic Resonance Spectroscopy to Monitor Therapy</b> .....	1464
Brian Ross, FRCS(Eng), London; James T. Helsper, MD, Pasadena, Calif; I. Jane Cox, PhD, London; Ian R. Young, PhD, Wembley, England; Raymond Kempf, MD, Los Angeles; Andrew Makepeace, MRCP, Jacqueline Pennock, BSc, London Tumor relapse or regression could be determined by magnetic resonance spectroscopy.	
<b>Laminin Expression in Colorectal Carcinomas Varying in Degree of Differentiation</b> .....	1470
George W. Daneker, Jr, MD; Arthur M. Mercurio, PhD; Lisa Guerra; Barbara Wolf, MD; Ronald R. Salem, FRCS; Darius J. Bagli, MD; Glenn D. Steele, Jr, MD, PhD, Boston The morphology and biochemistry of basement membranes are related to the degree of tumor differentiation.	
<b>Studies on Protein Kinase C and Colon Carcinogenesis</b> .....	1475
Jose G. Guillem, MD; Catherine A. O'Brian, PhD; Cheryl J. Fitzer; Mark D. Johnson, PhD; Kenneth A. Forde, MD; Paul LoGerfo, MD; I. Bernard Weinstein, MD, New York Bile salts induced a translocation of protein kinase C activity, which may relate to bile acid-mediated colon cancer.	
<b>Surgical Treatment of Spinal Chordomas</b> .....	1479
Narayan Sundaresan, MD; Andrew G. Huvos, MD; George Krol, MD; Joseph M. Lane, MD; Murray Brennan, MD, New York Computed tomography helps in the diagnosis of smaller and more resectable tumors.	
<b>Salvage Immunotherapy of Malignant Glioma</b> .....	1483
Marylou Ingram, MD; Skip Jacques, MD; Donald B. Freshwater, MD; Geza B. Techy, PhD; C. Hunter Sheldon, MD; James T. Helsper, MD, Pasadena, Calif Fifty patients with recurrent or residual malignant glioma responded initially to adoptive immunotherapy.	
<b>INDEX TO VOLUME 121</b> .....	1487

## REGULAR DEPARTMENTS

Instructions for Authors ..... See November 1987 Issue, p 1233

Index to Advertisers ..... 1362