

**PAPERS READ BEFORE THE 94TH ANNUAL MEETING OF THE
WESTERN SURGICAL ASSOCIATION, DEARBORN, MICH, NOV 16-19, 1986—Part I**

- Humor and the Surgeon: Presidential Address** 519
R. Dale Liechty, MD, Denver
"All of us must indulge in a few small follies if we are
to make reality bearable."—Proust
- Impact of a Trauma System on Outcome of Severely Injured Patients** 523
Steven R. Shackford, MD; Robert C. Mackersie, MD; David B. Hoyt, MD; William G. Baxt, MD; A. Brent Eastman, MD;
Fred N. Hammill, MD; F. Barry Knotts, MD; Richard W. Virgilio, MD, San Diego
Coordinated prehospital and hospital care of trauma victims
produced good survival results.
- Gastric Emptying and Small Bowel Transit of Solid Food
After Pylorus-Preserving Pancreaticoduodenectomy** 528
Marco G. Patti, MD; Carlos A. Pellegrini, MD; Lawrence W. Way, MD, San Francisco
Pylorus-preserving pancreaticoduodenectomy was associated
with normal gastric emptying.
- Criteria for the Selection of 'Early' Carcinomas of the Rectum:
Are They Valid?** 533
Jane C. Nelson, MD; Abdullatif N. Nimr, MD; Neil R. Thomford, MD, Toledo
Lymph node metastases were frequently present in histologically well-differentiated
rectal carcinomas less than 4 cm in size.
- Pelvic and Sacropelvic Exenteration for Locally
Advanced or Recurrent Anorectal Cancer** 537
Nathan W. Pearlman, MD; Robert E. Donohue, MD; Gregory V. Steigmann, MD;
Dennis J. Ahnen, MD; Scott M. Sedlacek, MD; Thomas J. Braun, MD, Denver
Radiation and exenteration were helpful in the treatment
of locally advanced pelvic anorectal carcinoma.
- Vitelline Duct Anomalies: Experience With 217 Childhood Cases** 542
Dennis W. Vane, MD; Karen W. West, MD; Jay L. Grosfeld, MD, Indianapolis
Meckel's diverticula produce many complications in infants and children, which suggests
that incidentally discovered diverticuli should be resected.
- Long-term Follow-up for Treatment of
Complicated Chronic Reflux Esophagitis** 548
Siroos S. Shirazi, MD; Konrad Schulze, MD; Robert T. Soper, MD, Iowa City
A short, loose Nissen fundoplication produced long-term cure of reflux esophagitis.
- Perforated Gastric Ulcers:
A Plea for Management by Primary Gastric Resection** 555
Gregory S. McGee, MD, John L. Sawyers, MD, Nashville, Tenn
A reduction in mortality and complications occurred when
primary resection rather than patch closure was performed.

The ARCHIVES OF SURGERY (ISSN 0004-0010) is published monthly by the American Medical Association, 535 N Dearborn St, Chicago, IL 60610, and is an official publication of the Association. Second-class postage paid at Chicago, IL 60610 and at additional mailing office.

SUBSCRIPTION RATES—The subscription rates for the Archives of Surgery are as follows: for members of the AMA, \$1.50 included in the annual membership dues; for nonmembers, \$38 for one year, \$68 for two years in the United States and US possessions; all other countries, one year, \$50; two years, \$92 (add \$15 surcharge to each one-year subscription, \$30 for two years for air delivery to the United Kingdom, Europe, Bangladesh, India, Nepal, or Sri Lanka). Special yearly rates to residents and medical students in the United States and US possessions, one year, \$19; two years, \$34. Rates for subscriptions for delivery to Australia, Japan, New Zealand, or South Korea are available through respective exclusive agents. Address all

subscription communications to the American Medical Association, Circulation and Fulfillment Division, 535 N Dearborn St, Chicago, IL 60610. Phone: (312) 280-7154.

CHANGE OF ADDRESS—POSTMASTER, send all address changes to Caryl L. Wertheimer, Director of Fulfillment, 535 N Dearborn St, Chicago, IL 60610. Notification of address change must be made at least six weeks in advance, include both old and new addresses, a recent mailing label, and your new zip code.

ADVERTISING PRINCIPLES: Each advertisement in this issue has been reviewed and complies with the principles governing advertising in AMA scientific publications. A copy of these principles is available on request. The appearance of advertising in AMA publications is not an AMA guarantee or endorsement of the product or the claims made for the product by the manufacturer.

Presentation and Perioperative Management of Arterioportal Fistulas 563
 William E. Strodel, MD; Frederic E. Eckhauser, MD; John H. Lemmer, MD;
 Walter M. Whitehouse, Jr, MD; David M. Williams, MD, Ann Arbor, Mich
 The development of an abdominal bruit in a patient who has had penetrating abdominal
 trauma suggests an arteriovenous fistula.

Pancreatitis After Biliary Tract Surgery 575
 Anthony Vernava, MD; Charles Andrus, MD; Virginia M. Herrmann, MD; Donald L. Kaminski, MD, St Louis
 Postoperative pancreatitis occurred in 8.4% of patients undergoing common duct exploration.

Late Results of Peripheral Vascular Surgery in Patients 80 Years of Age and Older 581
 Thomas H. Cogbill, MD; Jeffrey Landercasper, MD; Pamela J. Strutt, RN; Adolf L. Gundersen, MD, La Crosse, Wis
 Vascular reconstruction in the elderly was performed with
 low mortality and good results.

Duplex B-Mode Imaging for the Diagnosis of Deep Venous Thrombosis 587
 Mark Langsfeld, MD; Falls B. Hershey, MD; Lynn Thorpe, RN; Arthur I. Auer, MD;
 H. Bradley Binnington, MD; Joseph J. Hurley, MD; John J. Woods, MD, St Louis
 Duplex B-mode imaging and Doppler ultrasound effectively
 diagnosed lower-extremity deep-vein thrombosis.

**Mandatory vs Selective Exploration for Penetrating Neck Trauma:
 A Prospective Assessment** 592
 Joseph P. Meyer, MD; John A. Barrett, MD; James J. Schuler, MD; D. Preston Flanigan, MD, Chicago
 Clinical assessment alone is inadequate to predict major neck injuries.

Technical Defects as a Cause of Early Graft Failure After Femorodistal Bypass 599
 Larry L. Stept, MD; William R. Flinn, MD; Walter J. McCarthy III, MD;
 Stephen T. Bartlett, MD; John J. Bergan, MD; James S. T. Yao, MD, PhD, Chicago
 Technical problems were not the most common cause of early
 occlusion of vascular reconstructions.

**Diagnostic Pericardial Window:
 A Safe Alternative to Exploratory Thoracotomy for Suspected Heart Injuries** 605
 Frank B. Miller, MD; Sheldon J. Bond, MD; Charles R. Shumate, MD;
 Hiram C. Polk, Jr, MD; J. David Richardson, MD, Louisville
 Diagnosis of cardiac injury in patients with equivocal signs
 can be made quickly by a pericardial window.

Jejunioileal Bypass: A Legacy of Late Complications 610
 John R. Kirkpatrick, MD, Pittsburgh
 Reversal of all intestinal bypass operations performed
 for obesity is recommended.

**Neonatal Diaphragmatic Hernia:
 An Improving Outlook With Extracorporeal Membrane Oxygenation** 615
 Thomas R. Weber, MD; Robert H. Connors, MD; D. Glenn Pennington, MD; Susan Westfall, MD;
 William Keenan, MD; Suresh Kotagal, MD; J. Eugene Lewis, MD, St Louis
 Extracorporeal membrane oxygenation helps to improve survival
 in newborns with congenital diaphragmatic hernia.

REGULAR DEPARTMENTS

Correspondence and Brief Communications 619
Single-Valve vs Double-Valve Denver Peritoneovenous Shunts for Ascitic Fluid John B. Newkirk, DSc, Evergreen, Colo . . .
Subcutaneous Emphysema and Cocaine: A Rare Case Nagui Khouzam, MD, Winter Garden, Fla; Lenworth M. Jacobs, MD,
 Robert J. Schwartz, MD, Hartford, Conn . . . **Noninvasive Detection of Gallstone Acute Pancreatitis** Richard M. Gore, MD,
 Robert L. Vogelzang, MD, Chicago

Calendar 620

Ad Index 508

Instructions for Authors 517