

ORIGINAL ARTICLES

- Granulocyte Chemiluminescence in Patients With Postoperative Infections** 17
Matti Salo, MD; Juha Perttilä, MD; Olli-Pekka Lehtonen, MD, Turku, Finland
Oxygen-dependent microbicidal responses of granulocytes were depressed in patients with postoperative infections.
- Acquired Immunodeficiency Syndrome and Directed Blood Donations: A Dilemma for American Medicine** 23
Margot S. Kruskall, MD, Boston, Joel Umlas, MD, Cambridge, Mass
Directed donor programs have many pitfalls without proved medical value.
- Diagnosis and Operative Indications for Polypoid Lesions of the Gallbladder** 26
Akitoshi Koga, MD; Kiyooki Watanabe, MD; Tokihiko Fukuyama, MD;
Satoshi Takiguchi, MD; Fumio Nakayama, MD, Fukuoka, Japan
The malignant potential of gallbladder polyps is related to size.
- The Prevention of Pulmonary Complications After Upper Abdominal Surgery in Patients With Noncompromised Pulmonary Status** 30
Jan A. Roukema, MD; Eric J. Carol, MD; Jan G. Prins, MD, Eindhoven, the Netherlands
Preoperative and postoperative breathing exercises were associated with fewer pulmonary problems.
- α -Adrenergic Blocker for Posthernioplasty Urinary Retention: Prevention and Treatment** 35
Gideon Goldman, MD; Amos Leviav, MD; Amos Mazor, MD; Hanoeh Kashtan, MD; David Aladgem, MD;
Alexander Greenstein, MD; Theodor Wiznitzer, MD, Tel Aviv, Israel
Patients who received phenoxybenzamine orally before and after hernia repair did not develop urinary retention.
- The Role of Fine-Needle Aspiration in the Management of Solid Breast Masses** 37
Bruce Adye, MD; Philip C. Jolly, MD; Donald E. Bauermeister, MD, Seattle
The accuracy rate of cytologic evaluation of needle aspirates of breast masses is 80%.
- To Drain or Not to Drain in Thyroid Surgery: A Controlled Clinical Study** 40
Olle Wihlborg, MD; Lars Bergljung, MD, PhD; Hans Mårtensson, MD, PhD, Helsingborg, Sweden
There was no advantage to routine drainage in uncomplicated thyroid surgery.
- Intestinal Obstruction in Patients With Ovarian Cancer: Variables Associated With Surgical Complications and Survival** 42
Daniel L. Clarke-Pearson, MD; E. R. DeLong, PhD; Neeoo Chin, MD; Reed Rice, MD; William T. Creasman, MD, Durham, NC
Methods are provided to help select patients who might benefit from correction of obstruction.
- Correlation of Presence of Granulomas With Clinical and Immunologic Variables in Crohn's Disease** 46
Tomas M. Heimann, MD; Felice Miller; Giorgio Martinelli, PhD; Arnold Szporn, MD;
Adrian J. Greenstein, MD; Arthur H. Aufses, Jr, MD, New York
The presence of granulomas in patients with Crohn's disease was associated with high recurrence rates.
- Pain Insensitivity in Schizophrenic Patients: A Surgical Dilemma** 49
Linda K. Bickerstaff, MD; Stanley C. Harris, MD; Randi S. Leggett, MD; Keong-Chye Cheah, MD, Tacoma, Wash
The communication problems of mentally ill patients necessitate singularly insightful preoperative and postoperative care.
- Intramural Hematoma of the Duodenum** 54
Theodore C. Jewett, Jr, MD; Vincent Caldarola, MD; Melvyn P. Karp, MD; James E. Allen, MD; Donald R. Cooney, MD, Buffalo
Obstruction from a hematoma almost always subsides without an operation.
- Results of a Mass Screening Program for Colorectal Cancer** 63
Mark P. Miller, MD, Thomas V. Stanley, MD, Memphis
Stool guaiac slide kits returned by 23 000 persons identified 26 patients with colorectal cancers and 67 with polyps.

The ARCHIVES OF SURGERY (ISSN 0004-0010) is published monthly by the American Medical Association, 535 N Dearborn St, Chicago, IL 60610, and is an official publication of the Association. Second-class postage paid at Chicago, IL 60610 and at additional mailing office.

SUBSCRIPTION RATES—The subscription rates for the Archives of Surgery are as follows: \$40 for one year, \$70 for two years in the United States and US possessions; all other countries, one year, \$52; two years, \$96 (add \$15 surcharge for each one-year subscription, \$30 for two years for air delivery to the United Kingdom, Europe, Bangladesh, India, Nepal, or Sri Lanka). Rates for subscriptions for delivery to Australia, Japan, New Zealand, or South Korea are available through respective agents. **Special rates for residents and medical students in the United States and US possessions are available.** Address inquiries to the American Medical

Association, Circulation and Fulfillment Division, 535 N Dearborn St, Chicago, IL 60610. Phone: (312) 280-7168.

CHANGE OF ADDRESS—POSTMASTER, send all address changes to Caryl L. Wertheimer, Director of Fulfillment, 535 N Dearborn St, Chicago, IL 60610. Notification of address change must be made at least six weeks in advance, include both old and new addresses, a recent mailing label, and your new zip code.

ADVERTISING PRINCIPLES: Each advertisement in this issue has been reviewed and complies with the principles governing advertising in AMA scientific publications. A copy of these principles is available on request. The appearance of advertising in AMA publications is not an AMA guarantee or endorsement of the product or the claims made for the product by the manufacturer.

CONTENTS—Continued

Delayed Muscular Rigidity and Respiratory Depression Following Fentanyl Anesthesia	66
Joseph M. Klausner, MD, Tel Aviv, Israel; Joseph Caspi, MD, Hacarmel, Israel; Shlomo Lelcuk, MD, Aaron Khazam, MD, Tel Aviv, Israel; Gideon Marin, MD, Hacarmel, Israel; Herbert B. Hechtman, MD, Boston; Roland R. Rozin, MD, Tel Aviv, Israel Respiratory distress three to five hours after operation with fentanyl anesthesia may be a delayed toxic phenomenon.	
Diagnosis Related Groups and the Transfer of General Surgical Patients Between Hospitals	68
Eric Muñoz, MD, MBA; Richard Soldano, MPH; Howard Gross, MD, MS; Donald Chalfin, MD; Katherine Mulloy; Leslie Wise, MD, New Hyde Park, NY The data support the hypothesis that DRG reimbursement will provide a financial disincentive for teaching hospitals to accept surgical transfer patients. Invited Editorial Comment by Ira M. Rutkow, MD, MPH, DPH, Freehold, NJ	
Closure of Abdominal Wounds With Polydioxanone	72
David J. Schoetz, Jr, MD; John A. Collier, MD; Malcolm C. Veidenheimer, MD, Burlington, Mass Polydioxanone was safe and effective for closure of abdominal wounds.	

CLINICAL NOTES

Pseudohyperparathyroidism Secondary to Gigantic Mammary Hypertrophy	80
Jon A. Van Heerden, MB,ChB (Cape Town), FRCS(C); Hossein Gharib, MD; Ian T. Jackson, MB,ChB, FRCS, Rochester, Minn Gigantic mammary hypertrophy was associated with severe hypercalcemia that responded to mastectomies.	
Isolated Crohn's Disease of the Appendix: Two Case Reports and a Review of the Literature	85
Vincent W. Vanek, MD; George Spiritos, MD; Mounir Awad, MD; Narendra Badjatia, MD; Donald Bernat, MD, Youngstown, Ohio A low recurrence rate is associated with appendectomy for Crohn's disease isolated to the appendix.	
Fragmentation of Biliary Tract Stones by Lithotripsy Using Local Anesthesia	91
Bruce P. Brown, MD; Stefan A. Loening, MD; Fred C. Johlin, MD; Merril T. Dayton, MD; James W. Maher, MD, Iowa City Fragmentation of stones was accomplished in two patients. Editorial Comment by David L. Nahrwold, MD, Chicago	

BRIEF CLINICAL NOTES

Adhesion-Related Variceal Hemorrhage Following Sclerosis of Esophageal Varices	94
Robert D. Fry, MD; Keith C. Fischer, MD; Noah Susman, MD; Burton A. Shatz, MD; Bernard Hulbert, MD, St Louis Varices forming in intestinal adhesions are an unusual cause of lower gastrointestinal tract hemorrhage.	
Villous Adenoma of the Common Bile Duct	96
Jonathan Saxe, MD; Charles Lucas, MD; Anna M. Ledgerwood, MD; Choichi Sugawa, MD, Detroit Villous adenoma of the distal common bile duct was treated by pancreaticoduodenectomy.	
Bran-Induced Small-Intestinal Obstruction in a Patient With No History of Abdominal Operation	98
James B. McClurken, MD, Ned Z. Carp, MD, Abington, Pa Eating two large bowls of bran cereal led to intestinal obstruction in this patient.	

TECHNIQUES

In Situ Grafting Made Easy: Modification of a Technique	101
George D. LeMaitre, MD, Michael J. Arakelian, MD, Bon Secours, Mass Use of a retrograde valvulotome and Doppler for locating tributaries followed by the distal anastomosis has been time-saving and effective.	
Ultrasonic Guided Percutaneous Transhepatic Bile Drainage for Cholangitis due to Intrahepatic Stones	106
Min-Ho Huang, MD, Changhua, Taiwan, Republic of China, Chen-Guo Ker, MD, Kaohsiung, Taiwan, Republic of China This technique helped control the emergency problem in 41 patients.	
Modification of the End-to-End Stapled Anastomosis Utilizing Intersecting Staple Lines	110
Karl A. LeBlanc, MD, Vincent R. Russo, MD, New Orleans An EEA staple row intersects a GIA staple line in this technique of colon anastomosis.	
Pleurectomy Through the Triangle of Auscultation for Treatment of Recurrent Pneumothorax in Younger Patients: Evaluation of 60 Consecutive Cases	113
Jamshid Nazarian, MD; G. Down, MBDS; O. J. Lau, FRCS, London This limited thoracotomy avoids muscle transection.	

SPECIAL ARTICLE

An Experiment in Surgical Education—the First International Exchange of Residents: The Letters of Halsted, Küttner, Heuer, and Landols	115
Ira M. Rutkow, MD, MPH, DrPH, Freehold, NJ, Karl Hempel, MD, Hamburg, West Germany Letters between Halsted of Hopkins and Küttner of Breslau describe the exchange of residents before World War I.	

REGULAR DEPARTMENTS

Correspondence and Brief Communications	122
Books	125
Calendar	127
Instructions for Authors	8
Index to Advertisers	128