

**PAPERS READ BEFORE THE SEVENTH ANNUAL MEETING OF THE
SURGICAL INFECTION SOCIETY, PHILADELPHIA,
MAY 11 TO MAY 12, 1987—PART I****Postresidency Fellowships: An Investment In the Future:**

- Presidential Address** 147
William S. Blakemore, MD, Birmingham, Ala

Treatment of Severe Intra-abdominal Sepsis and/or Necrotic Foci by an

- 'Open-Abdomen' Approach: Zipper and Zipper-Mesh Techniques** 152
J. L. Garcia-Sabrido, MD, PhD; J. M. Tallado, MD; N. V. Christou, MD, PhD;
J. R. Polo, MD, PhD; E. Valdecantos, MD, Madrid
Daily laparotomies in the ICU with epidural catheter anesthesia
and abdominal zippers seemed helpful.

The Comparative Clearance Rates of the Pleural and Peritoneal Cavities 157

- Constantine Mavroudis, MD; Mark A. Malangoni, MD; Sara L. Katzmark; Richard Montgomery;
C. Randolph Schrodt, MD; Hiram C. Polk, Jr, MD, Louisville
Clearance of bacteria from the pleural space was greater than
from the peritoneum in rats.

Inhibition of Cyclo-oxygenase Attenuates the Metabolic

- Response to Endotoxin in Humans** 162
Arthur Revhaug, MD; Hamish R. Michie, FRCS(E); James McK. Manson, FRCS(E); James M. Watters, MD;
Charles A. Dinarello, MD; Sheldon M. Wolff, MD; Douglas W. Wilmore, MD, Boston
The flu-like symptoms of fever, tachycardia, and increased metabolic rate and
stress hormone release following endotoxin injection were attenuated by ibuprofen.

Neutrophil Activation in Sepsis: The Relationship Between fMet-Leu-Phe Receptor

- Mobilization and Oxidative Activity** 171
Stephen D. Tennenberg, MD, Joseph S. Solomkin, MD, Cincinnati
Neutrophils from septic patients had surface receptor expression
that did not correlate with functional activity.

The Microbiology of Neonatal Peritonitis 176

- Daniel L. Mollitt, MD; Joseph J. Tepas III, MD; James L. Talbert, MD, Jacksonville, Fla
Classic enteric antibiotic coverage may be inappropriate and ineffective in
the newborn with peritonitis.

Allosensitized Helper and Cytotoxic T-lymphocyte Clones

- Differentially Modulate Endotoxin-Stimulated Macrophage Function** 182
Mark L. Jordan, MD; Ann Carlson; Rosemary A. Hoffman; Richard L. Simmons, MD, Minneapolis
Helper and cytotoxic T cells altered the production of IL-1 and PGE₂ by
endotoxin-activated macrophages.

The ARCHIVES OF SURGERY (ISSN 0004-0010) is published monthly by the American Medical Association, 535 N Dearborn St, Chicago, IL 60610, and is an official publication of the Association. Second-class postage paid at Chicago, IL 60610 and at additional mailing office.

SUBSCRIPTION RATES—The subscription rates for the Archives of Surgery are as follows: \$40 for one year, \$70 for two years in the United States and US possessions; all other countries, one year, \$52; two years, \$96 (add \$15 surcharge for each one-year subscription, \$30 for two years for air delivery to the United Kingdom, Europe, Bangladesh, India, Nepal, or Sri Lanka). Rates for subscriptions for delivery to Australia, Japan, New Zealand, or South Korea are available through respective agents. *Special rates for residents and medical students in the United States and US possessions are available.* Address inquiries to the American Medical

Association, Circulation and Fulfillment Division, 535 N Dearborn St, Chicago, IL 60610. Phone: (312) 280-7168.

CHANGE OF ADDRESS—POSTMASTER, send all address changes to Caryl L. Wertheimer, Director of Fulfillment, 535 N Dearborn St, Chicago, IL 60610. Notification of address change must be made at least six weeks in advance, include both old and new addresses, a recent mailing label, and your new zip code.

ADVERTISING PRINCIPLES: Each advertisement in this issue has been reviewed and complies with the principles governing advertising in AMA scientific publications. A copy of these principles is available on request. The appearance of advertising in AMA publications is not an AMA guarantee or endorsement of the product or the claims made for the product by the manufacturer.

CONTENTS—Continued

Terminal Complement Complexes and Anaphylatoxins in Septic and Ischemic Patients	188
Mats Heideman, MD, PhD; Birgitta Norder-Hansson; Anders Bengtson, MD, PhD, Gothenburg, Sweden; Tom Eirik Mollnes, MD, PhD, Oslo	
Evidence for complement activation in patients with sepsis or ischemia returned to normal with successful therapy.	
Candida: A Decreasing Problem for the Burned Patient?	194
Baiba J. Grube, MD; Janet A. Marvin, RN, MN; David M. Heimbach, MD, Seattle	
<i>Candida</i> septicemia was not a major cause of morbidity or mortality in this burn unit.	
Coagulation Augments Neutrophil C3b Receptors via Formation of a Protein(s) Unrelated to Fibrinolysis or C5 Activation	199
Julio Garcia-Aguilar, MD; Marc E. Lanser, MD; Glenn E. Brown, PhD, Boston	
Increased expression of complement receptors in cell membranes was produced by platelet activation.	
The Effect of the Immunomodulator RU 41 740 (Biostim) on the Specific and Nonspecific Immunosuppression Induced by Thermal Injury or Protein Deprivation	207
Nicolas V. Christou, MD, PhD; Ihor Zakaluzny, MD; John C. Marshall, MD; Carl W. Nohr, MD, Montreal	
A macrophage stimulator produced the opposite results with a burn injury as compared with protein deprivation.	
Red Blood Cell Deformability in Human and Experimental Sepsis	217
Thelma C. Hurd, MD; Kumar S. Dasmahapatra, MD; Benjamin F. Rush, Jr, MD; George W. Machiedo, MD, East Orange, NJ	
Red blood cells from septic patients had decreased deformability, as did cells from septic or endotoxemic rats.	
Evidence That Cathepsin B Contributes to Skeletal Muscle Protein Breakdown During Sepsis	221
Robert P. Hummel III; J. Howard James; Brad W. Warner, MD; Per-Olof Hasselgren, MD, PhD; Josef E. Fischer, MD, Cincinnati	
Lysosomal protease activity and protein degradation rate were both increased in muscles from septic animals.	
APACHE II Score and Abdominal Sepsis: A Prospective Study	225
John M. A. Bohnen, MD, FRCS(C); Robert A. Mustard, MD, FRCS(C); Susan E. Oxholm, RN; B. Diana Schouten, RN, Toronto	
The Acute Physiology and Chronic Health Evaluation (APACHE) score and steroid use were associated with increased mortality.	
Fibrin Sealant in High-Risk Colonic Anastomoses	230
Kimberly A. Houston, MSc, Ori D. Rotstein, MD, Toronto	
Fibrin sealant had no effect on colon anastomoses in rats.	
Effect of Blood Transfusion on Antigen Presentation Function and on Interleukin 2 Generation	235
Rabie N. Stephan, MD; John M. Kisala, MD; Richard E. Dean, MD, East Lansing, Mich; Alexander S. Geha, MD, Cleveland; Irshad H. Chaudry, PhD, East Lansing, Mich	
The ability of mouse splenocytes to generate interleukin 2 decreased with a single blood transfusion.	
Enhancement of Mitochondrial Function in Sepsis	241
Konrad L. Dawson, MD, Evan R. Geller, MD, Detroit; John R. Kirkpatrick, MD, Pittsburgh	
<i>Escherichia coli</i> endotoxin was associated with increased mitochondrial function in skeletal muscle, heart, and liver.	
Soft-Tissue Infections Caused by Halophilic Marine Vibrios	245
Richard J. Howard, MD, Gainesville, Fla, Spencer Lieb, MPH, Tallahassee, Fla	
Cellulitis in a patient who has been near ocean water should suggest this possibility.	
PANEL DISCUSSION—SURGICAL INFECTION SOCIETY 1987	
Does Infection Control Control Infection?	250
Robert E. Condon, MD, Milwaukee; Robert W. Haley, MD, Dallas; James T. Lee, Jr, MD, PhD, Minneapolis; Jonathan L. Meakins, MD, Montreal	
Certainly, when properly done!	
REGULAR DEPARTMENTS	
Correspondence and Brief Communications	258
Books	264
Calendar	267
Instructions for Authors	See January 1988 issue, p 8
Index to Advertisers	262