

## EDITORIALS

- Hypothesis: Sepsis, Multiple Systems Organ Failure, and the Macrophage** ..... 285  
John R. Border, MD, Buffalo

**PAPERS READ BEFORE THE ANNUAL MEETING OF THE SURGICAL INFECTION SOCIETY  
PHILADELPHIA, MAY 11 TO MAY 12, 1987—PART II**

- Alteration of Monocyte Function Following Major Injury** ..... 287  
Eugen Faist, MD; Angelika Mewes, MD; Theodor Strasser, MD, Munich; Alfred Waiz, MD, Bern, Switzerland;  
Sefik Alkan, MD, Basel, Switzerland; Christopher Baker, MD, New Haven, Conn; Wolfgang Ertel, MD; George Heberer, MD, Munich  
Negative regulatory macrophage function may be the event initiating posttraumatic immunosuppression.
- Mechanisms of Altered Monocyte Prostaglandin E<sub>2</sub>  
Production in Severely Injured Patients** ..... 293  
Carol L. Miller-Graziano, PhD; Mitchell Fink, MD; Jia Yan Wu, PhD; Gyongyi Szabo, MD; Karen Kodys, Worcester, Mass  
Elevated monocyte prostaglandin E<sub>2</sub> synthesis due to a shift in monocyte subset ratios may  
contribute to immunosuppression.
- Hydrogen Peroxide-Mediated Inhibition of T-Cell Response  
to Mitogens Is a Result of Direct Action on T Cells** ..... 300  
Deborah A. Patterson, MA; Robert Rapoport; Mark A. K. Patterson, PhD; Brian M. Freed, MA; Neil Lempert, MD, Albany, NY  
Hydrogen peroxide from inflammatory cells may suppress nearby T lymphocytes.
- Generation of an Anti-Interleukin 2 Factor in Healing Wounds** ..... 305  
Robert J. Breslin, MD; Adrian Barbul, MD; Thomas S. Kupper, MD;  
John P. Knud-Hansen, MD; Hannah L. Wasserkrug; Gershon Efron, MD, Baltimore  
Wound fluid contains a factor(s) that impairs lymphocyte  
response to interleukin 2 but enhances fibroblast proliferation.
- The Microbiology of Multiple Organ Failure:  
The Proximal Gastrointestinal Tract as an Occult Reservoir of Pathogens** ..... 309  
John C. Marshall, MD, FRCSC; Nicholas V. Christou, MD, PhD, FRCSC, FACS;  
Ruth Horn, MD, FRCPC; Jonathan L. Meakins, MD, DSc, FRCSC, FACS, Montreal  
Pathogens that dominate in patients with multiple organ failure also are found in the upper gastrointestinal tract.
- Systemic Complement Activation Produces Hemodynamic  
Changes Characteristic of Sepsis** ..... 316  
William J. Schirmer, MD; James M. Schirmer; George B. Naff, MD; Donald E. Fry, MD, Cleveland  
Zymosan injection to produce complement activation also  
stimulated the hemodynamic changes of hyperdynamic sepsis.
- Prospective Study Comparing Imipenem-Cilastatin With Clindamycin and Gentamicin  
for the Treatment of Serious Surgical Infections** ..... 322  
Alan W. Hackford, MD; Francis P. Tally, MD; Randolph B. Reinhold, MD; Michael Barza, MD; Sherwood L. Gorbach, MD, Boston  
Single-agent therapy with imipenem was as effective as combined therapy with the other agents.
- Splenectomy Alters Kupffer Cell Response to Endotoxin** ..... 327  
Timothy R. Billiar, MD; Michael A. West, MD; Bonnie J. Hyland; Richard L. Simmons, MD, Minneapolis  
Decreased sensitivity of Kupffer cells after splenectomy may be due to loss of priming by splenic lymphokines.
- Duration of Preventive Antibiotic Administration for Open Extremity Fractures** ..... 333  
E. Patchen Dellinger, MD, Seattle; Ellis S. Caplan, MD, Baltimore; Lance D. Weaver, MD, Los Angeles; Margaret J. Wertz, MN, RN,  
Beth M. Droppert, RN, Seattle; Nancy Hoyt, MACIC; Robert Brumback, MD; Andrew Burgess, MD; Attila Poka, MD, Baltimore;  
Stephen K. Benirschke, MD, E. Stan Lennard, MD, Seattle; Sister Mary Ann Lou, MD, Los Angeles  
One day of antibiotic therapy was as effective as five days for open fractures.

The ARCHIVES OF SURGERY (ISSN 0004-0010) is published monthly by the American Medical Association, 535 N Dearborn St, Chicago, IL 60610, and is an official publication of the Association. Second-class postage paid at Chicago, IL 60610 and at additional mailing office.

**SUBSCRIPTION RATES**—The subscription rates for the Archives of Surgery are as follows: \$40 for one year, \$70 for two years in the United States and US possessions; all other countries, one year, \$52; two years, \$96 (add \$15 surcharge for each one-year subscription, \$30 for two years for air delivery to the United Kingdom, Europe, Bangladesh, India, Nepal, or Sri Lanka). Rates for subscriptions for delivery to Australia, Japan, New Zealand, or South Korea are available through respective agents. **Special rates for residents and medical students in the United States and US possessions are available.** Address inquiries to the American Medical

Association, Circulation and Fulfillment Division, 535 N Dearborn St, Chicago, IL 60610. Phone: (312) 280-7168.

**CHANGE OF ADDRESS—POSTMASTER**, send all address changes to Caryl L. Wertheimer, Director of Fulfillment, 535 N Dearborn St, Chicago, IL 60610. Notification of address change must be made at least six weeks in advance, include both old and new addresses, a recent mailing label, and your new zip code.

**ADVERTISING PRINCIPLES:** Each advertisement in this issue has been reviewed and complies with the principles governing advertising in AMA scientific publications. A copy of these principles is available on request. The appearance of advertising in AMA publications is not an AMA guarantee or endorsement of the product or the claims made for the product by the manufacturer.

**CONTENTS—Continued**

**Direct Effects of Endotoxin on Hepatocytes: Synthesis of a Specific Secretory Protein** ..... 340  
John E. Mazuski, MD; Jeffrey L. Platt, MD; Michael A. West, MD;  
Richard L. Simmons, MD; Howard C. Towle, PhD; Frank B. Cerra, MD, Minneapolis  
Synthesis of a protein by hepatocytes was stimulated by glucocorticoids, interleukin 1, and endotoxin.

**Pulmonary Microvascular Changes Following Fluid Resuscitation in an Ovine Model of Endotoxemia** ..... 345  
Hans J. Lubbesmeyer, MD; Ryozo Kimura, MD; James P. Maguire, MD; Mike Irei, MD;  
Lillian D. Traber, RN; Daniel L. Traber, PhD; David N. Herndon, MD, Galveston, Tex  
Augmented fluid resuscitation during endotoxemia increased pulmonary microvascular pressure and fluid flux but did not decrease arterial  $PO_2$ .

**SPECIAL ARTICLE**

**Injury to Segmental Bile Ducts: A Reappraisal** ..... 351  
Nicholas S. Hadjis, FRCS, Leslie H. Blumgart, FRCS, London  
Should a segmental bile duct injury be ligated or reconstructed?  
Invited Editorial Comment by John W. Braasch, MD, Burlington, Mass

**ORIGINAL ARTICLES**

**Computer-Derived Equations for Predicting Survival Postoperatively: Their Usefulness and Limitations** ..... 354  
Eric B. Rypins, MD; Farrukh Khan, MD; Dawn Collins-Irby, RN; I. James Sarfeh, MD, Long Beach, Calif;  
James T. Ashurst, PhD, Irvine, Calif; Edward A. Stemmer, MD, Long Beach, Calif  
Mortality after GI tract perforation could be predicted by age of the patient, presence of pulmonary disease, and preoperative shock.  
Invited Editorial Comment by John H. Siegel, MD, Baltimore

**Infectious Complications in Liver Transplantation** ..... 360  
John O. Colonna II, MD; Drew J. Winston, MD; Judith E. Brill, MD; Leonard I. Goldstein, MD; Marilyn P. Hoff, RN;  
Jonathan R. Hiatt, MD; William Quinones-Baldrich, MD; Kenneth P. Ramming, MD; Ronald W. Busuttil, MD, PhD, Los Angeles  
Infections developed in two thirds of patients receiving liver transplants.

**Septicemia From Biliary Tract Infection** ..... 366  
Yardena Siegman-Igra, MD; David Schwartz, MSc; Nissim Konforti, MSc;  
Chaim Perluk, MD; Roland R. Rozin, MD, Tel-Aviv, Israel  
The most common microorganisms were *Escherichia coli*, *Klebsiella*, *Streptococcus*, and *Proteus*.

**Splenectomy for Thrombocytopenia due to Secondary Hypersplenism** ..... 369  
William W. Coon, MD, Ann Arbor  
Patients with thrombocytopenia associated with chronic lymphocytic and hairy-cell leukemia and stage IV lymphoma may benefit from splenectomy.

**CLINICAL NOTES**

**Gastritis Cystica Polyposa** ..... 372  
Ahmet M. Özeng, MD; Şevket Ruacan, MD; Ömer Aran, MD, Ankara, Turkey  
Mucosal cystic hyperplasia at gastrojejunostomy sites produced bleeding and obstruction.

**Emphysematous Pyelonephritis in a Xanthogranulomatous Kidney: An Unusual Case of Pneumoperitoneum** ..... 377  
Lorrie A. Langdale, MD; Charles L. Rice, MD; Nancy Brown, MD, Chicago  
This patient developed increasing abdominal pain and air and pus in the peritoneal cavity due to pyelonephritis.

**Toothbrush Swallowing** ..... 382  
Allan D. Kirk; Bert A. Bowers, MD; Joseph A. Moylan, MD; William C. Meyers, MD, Durham, NC  
A toothbrush that is swallowed will not pass spontaneously and should be removed by endoscopy.

**TECHNIQUES**

**Effective Technique of Sternum Closure in High-Risk Patients** ..... 386  
Gideon Goldman, MD; Roberto Nestel, MD; Eitan Snir, MD; Bernardo Vidne, MD, Tel Aviv, Israel  
There were no dehiscences or mediastinitis in 100 high-risk patients who had sternal closures by a combination of mattress sutures and single sutures.

**Total Pharyngeal Reconstruction Using a Pectoralis Major Myocutaneous Tunnel** ..... 391  
William A. Maddox, MD; Christopher J. O'Brien, MS, FRACS; Lev Bragg, MD; Marshall M. Urist, MD, Birmingham, Ala  
A myocutaneous tunnel, rather than a tube, was used to reconstruct the cervical esophagus.

**REGULAR DEPARTMENTS**

**Correspondence and Brief Communications** ..... 394

**Books** ..... 395

**Calendar** ..... 396

**Instructions for Authors** ..... See January 1988 issue, p 8

**Index to Advertisers** ..... 390