

PAPERS READ BEFORE THE 68TH ANNUAL MEETING OF THE NEW ENGLAND SURGICAL SOCIETY, BRETTON WOODS, NH, SEPT 11 TO SEPT 13, 1987

The Many Faces of Surgery: Presidential Address	543
Joseph E. Murray, MD, Boston	
Beyond Transplantation: Third Annual Samuel Jason Mixter Lecture	545
Joseph P. Vacanti, MD, Boston	
Conservation of the Spleen With Distal Pancreatectomy	550
Andrew L. Warshaw, MD, Boston	
The distal pancreas can be mobilized and resected with preservation of a normal spleen if the short gastric vessels are preserved.	
Ripstein Procedure: Lahey Clinic Experience: 1963-1985	554
Patricia L. Roberts, MD; David J. Schoetz, Jr, MD; John A. Coller, MD; Malcolm C. Veidenheimer, MD, Burlington, Mass	
Recurrent prolapse developed in about 10% of patients.	
Safe Management of the Impossible Duodenum: Risk Avoidance in Surgery of Peptic Ulcer	558
Grant V. Rodkey, MD, Boston	
Partial antrectomy with intramural mucosal removal and gastroduodenostomy was safe and effective for the difficult duodenum.	
Surgical Treatment of Biliary Cystadenoma: A Report of 15 Cases	563
W. David Lewis, MD; Roger L. Jenkins, MD; Ricardo L. Rossi, MD, Boston; Lawrence Munson, MD, Stephen G. ReMine, MD, Burlington, Mass; Blake Cady, MD; John W. Braasch, MD; William V. McDermott, MD, Boston	
Complete excision of these unusual biliary tract lesions is recommended.	
Surgically Incurable Well-Differentiated Thyroid Carcinoma: Prognostic Factors and Results of Therapy	569
Ricardo L. Rossi, MD, Burlington, Mass; Blake Cady, MD, Boston; Mark L. Silverman, MD; Marvin S. Wool, MD; Steven G. ReMine, MD; Mary Beth Hodge, MD; Ferdinand A. Salzman, MD, Burlington, Mass	
Patients with surgically incurable but well-differentiated thyroid carcinoma should undergo resection of as much disease as possible.	
Predicting Hospital Charges for Trauma Care	579
Susan E. Pories, MD; Richard L. Gamelli, MD; Pamela Vacek, MS; Frank Harris, MS; Dean Lea, MS, Burlington, Vt	
Fair reimbursement to hospitals for trauma patients requires a modification of the DRG according to the Injury Severity Score and age of the patient.	
Primary Lung Cancer Surgery in Stage II and Stage III	583
Wilford B. Neptune, MD, Boston	
In the absence of contraindications, operative excision seems to offer the best possibility of survival in advanced lung cancer.	
Treatment of Crohn's Colitis: Segmental or Total Colectomy?	588
Walter E. Longo, MD; Garth H. Ballantyne, MD; C. Elton Cahow, MD, New Haven, Conn	
Recurrence is not related to the extent of resection, and segmental resection delays the need for a stoma and preserves functioning bowel.	
Ultra-High-Frequency Jet Ventilation in a Bronchopleural Fistula Model	591
Rocco Orlando III, MD; Eric H. Gluck, MD; Murray Cohen, MD; C. G. Mesologites, MD, Hartford, Conn	
Ultra-high-frequency ventilation was most effective in this animal bronchopleural fistula model.	
Persistent Acute Pancreatitis: A Variant Treated by Pancreatoduodenectomy	597
Peter L. Rutledge, MD, Andrew L. Warshaw, MD, Boston	
Acute pancreatitis that failed to subside over six to 12 weeks in five patients was treated by pancreatic resection.	

The ARCHIVES OF SURGERY (ISSN 0004-0010) is published monthly by the American Medical Association, 535 N Dearborn St, Chicago, IL 60610, and is an official publication of the Association. Second-class postage paid at Chicago, IL 60610 and at additional mailing office.

SUBSCRIPTION RATES—The subscription rates for the Archives of Surgery are as follows: \$40 for one year, \$70 for two years in the United States and US possessions; all other countries, one year, \$52; two years, \$96 (add \$15 surcharge for each one-year subscription, \$30 for two years for air delivery to the United Kingdom, Europe, Bangladesh, India, Nepal, or Sri Lanka). Rates for subscriptions for delivery to Australia, Japan, New Zealand, or South Korea are available through respective agents. *Special rates for residents and medical students in the United States and US possessions are available.* Address inquiries to the American Medical

Association, Circulation and Fulfillment Division, 535 N Dearborn St, Chicago, IL 60610. Phone: (312) 280-7168.

CHANGE OF ADDRESS—POSTMASTER, send all address changes to Caryl L. Wertheimer, Director of Fulfillment, 535 N Dearborn St, Chicago, IL 60610. Notification of address change must be made at least six weeks in advance, include both old and new addresses, a recent mailing label, and your new zip code.

ADVERTISING PRINCIPLES: Each advertisement in this issue has been reviewed and complies with the principles governing advertising in AMA scientific publications. A copy of these principles is available on request. The appearance of advertising in AMA publications is not an AMA guarantee or endorsement of the product or the claims made for the product by the manufacturer.

CONTENTS—Continued

Use of Oxygen Radical Scavengers on Autografted Pig Kidneys After Warm Ischemia and 48-Hour Perfusion Preservation	601
Peter J. Bosco, MD, Robert T. Schweizer, MD, Hartford, Conn Renal function was improved in kidneys treated with superoxide dismutase and catalase after normothermic ischemia and perfusion preservation.	
Effect of β-Adrenergic Blockade on the Growth Rate of Abdominal Aortic Aneurysms	606
Steven D. Leach, MD; Allan L. Toole, MD; Harold Stern, MD; Ralph W. DeNatale, MD; M. David Tilson, MD, New Haven, Conn Treatment by β -adrenergic blockade was associated with a lower growth rate of abdominal aortic aneurysms.	
Renal Failure and Advanced Atherosclerotic Lesions: Salvage by Vascular Reconstruction	610
James R. Elmore, MD; Ferris S. Ray, MD; Richard C. Dillihunt, MD; William E. Herbert, MD, Portland, Me Renal revascularization was helpful in 21 of 30 patients with renal insufficiency.	
Parathyroid Gland Exploration With Local Anesthesia in Elderly and High-Risk Patients	614
Ludwig J. Pyrtek, MD; Michael Belkin, MD; Stanley Bartus, MD; Robert Schweizer, MD, Hartford, Conn With good preoperative localization, the lateral approach using local anesthesia was safe and effective in these patients.	
Problems Associated With a Nissen Fundoplication Following Tracheoesophageal Fistula and Esophageal Atresia Repair	618
Michael R. Curci, MD, Albert W. Dibbins, MD, Portland, Me A fundoplication creates a mechanical obstruction in infants with esophageal motility problems associated with esophageal atresia-fistula.	
Combined Kidney and Pancreas Transplantation in Diabetics	621
A. Benedict Cosimi, MD; Hugh Auchincloss, Jr, MD; Francis L. Delmonico, MD; Leslie Fang, MD; David M. Nathan, MD; Nina Tolkoff-Rubin, MD; Robert H. Rubin, MD; Harold C. Yang, MD; Paul S. Russell, MD, Boston The success of pancreas transplantation in diabetics with advanced complications now approaches that of other organ allografts.	
Advances in the Diagnosis and Treatment of Pheochromocytoma	626
Robert J. Havlik, MD; C. Elton Cahow, MD; Barbara K. Kinder, MD, New Haven, Conn Epinephrine-norepinephrine fractionation, scanning technique, and selective α_1 antagonists have improved the diagnosis and therapy of pheochromocytomas.	
Pericardial Flap Prevents Sternal Wound Complications	636
William C. Nugent, MD; Elizabeth L. Maislen, RN; Gerald T. O'Connor, PhD, DSc; Charles A. S. Mairrin, MB, BS; Stephen K. Plume, MD, Hanover, NH The routine use of a pericardial flap interposed between the heart and sternum was associated with decreased mediastinitis and sternal dehiscence.	
ORIGINAL ARTICLES	
Protein Synthesis and Degradation in Skeletal Muscle From Septic Rats: Response to Leucine and α-Ketoisocaproic Acid	640
Per-Olof Hasselgren, MD, PhD; J. Howard James; Brad W. Warner, MD; Robert P. Hummel III; Josef E. Fischer, MD, Cincinnati Sepsis induces changes in skeletal muscle protein turnover that are resistant to the effects of leucine.	
Anaphylatoxin Formation in Sepsis	645
Anders Bengtson, MD, Mats Heideman, MD, PhD, Gothenburg, Sweden Persistent elevation of anaphylatoxins was strongly correlated with development of multiple organ failure.	
CLINICAL NOTES	
Septic Complications of Hemorrhoidal Banding	650
Gerardo Quevedo-Bonilla, MD; Anna M. Farkas, MD; Herand Abcarian, MD; Ernestine Hambrick, MD; Charles P. Orsay, MD, Chicago Rubber band ligation of hemorrhoids can lead to perianal abscess and systemic sepsis.	
TECHNIQUES	
The Scalp as a Donor Site in Burns	652
Edmond A. Zingaro, MD; Angelo Capozzi, MD; Vincent R. Pennisi, MD, San Francisco Complete hair regrowth has occurred in nearly all patients.	
A Method for Repair of Aortic Dissection Originating in the Transverse Arch Using Two Sutureless Intraluminal Prostheses	656
Eric Thompson, MD, Hartzell V. Schaff, MD, Rochester, Minn An intraluminal prosthesis facilitated the repair of acute dissection originating in the transverse aortic arch.	
Treatment of Liver Hemangiomas by Enucleation	659
Aydin Alper, MD; Orhan Ariogul, MD; Ali Emre, MD; Ali Uras, MD; Attila Okten, MD, Istanbul, Turkey Enucleation of hepatic hemangiomas simplifies surgical treatment.	
REGULAR DEPARTMENTS	
Correspondence and Brief Communications	662
Calendar	664
Instructions for Authors	536
Index to Advertisers	537