

ORIGINAL ARTICLES

- Breast Cancer in Women After Augmentation Mammoplasty** 681
Melvin J. Silverstein, MD; Neal Handel, MD; Parvis Gamagami, MD;
James R. Waisman, MD; Eugene D. Gierson, MD; Robert J. Rosser, MD;
Robert Steyskal, MD; William Colburn, MD, Van Nuys, Calif
Patients with augmentation mammoplasty who developed
cancer seemed to have more advanced disease.
Editorial Comment by LaSalle D. Leffall, Jr, MD, Washington, DC
- Neutrophil Intracellular Kill Following Thermal Injury: Different Bactericidal Capability
for Patients' Organisms and Laboratory Organisms** 686
Kevin T. Moran, MD; Maria M. Allo, MD; Thomas J. O'Reilly, MD; Diane F. Karadsheh; Andrew M. Munster, MD, Baltimore
Neutrophils from patients who died failed to kill their own as well as laboratory organisms.
- Simple Ligation vs Stump Inversion in Appendectomy** 689
David Street, MD; Balazs I. Bodai, MD; Leon J. Owens, MD; Daniel B. Moore, MD;
Charles B. Walton, MD; James W. Holcroft, MD, Davis, Calif
Adhesions requiring repeated operation for bowel obstruction occurred only
in patients who had stump inversion.
- A Review of the Histologic Changes in Vein-to-Artery Grafts, With Particular Reference to Intimal Hyperplasia** 691
Rodney J. Dilley, BScHons, PhD, John K. McGeachie, BSc, PhD, FDSRCS, Nedlands, Western Australia;
Francis J. Prendergast, MBBS, FRACS, Perth, Western Australia
Intimal hyperplasia developing in vein-to-artery grafts appears to be of smooth-muscle origin.
- Common Bile Duct Pressure in Patients With Common Bile Duct Stones
With or Without Acute Suppurative Cholangitis** 697
Attila Csendes, MD; Alfredo Sepúlveda, MD; Patricio Burdiles, MD; Italo Braghetto, MD;
Jorge Bastias, MD; Hans Schütte, MD; Juan C. Díaz, MD; Julio Yarmuch, MD; Fernando Maluenda, MD, Santiago, Chile
Increased common bile duct pressures and impacted stones were characteristic
of patients with acute suppurative cholangitis.
- Evaluation of Naloxone Therapy for *Escherichia coli* Sepsis in the Baboon** 700
Lerner B. Hinshaw, PhD; Linda T. Archer, PhD; Beverly K. Beller; Alvin C. K. Chang; Dayl J. Flournoy, PhD;
Richard B. Passey, PhD, Oklahoma City; Joseph B. Long, PhD, John W. Holaday, PhD, Washington, DC
The opioid antagonist, naloxone, had no therapeutic
benefit when evaluated in baboons with septic shock.
- Effects of Female Sex Hormones and Pregnancy on Gallbladder Prostaglandin Synthesis** 705
Eddie L. Hoover, MD; Bernard M. Jaffe, MD; Huelaine Webb, MD; David W. England, MB, ChB, Brooklyn, NY
A high-cholesterol diet increases rabbit gallbladder prostanoid formation,
but female sex hormones and pregnancy do not.
- Pneumoperitoneum and Its Association With Ruptured Abdominal Viscus** 709
Thomas G. Winek, MD; H. Steven Moseley, MD; Gordon Grout, MD; David Luallin, MD, Portland, Ore
Of 77 patients with free air, by x-ray only four did not have a perforated viscus.
- Moxalactam vs Tobramycin-Ciindamycin: A Randomized Trial in Secondary Peritonitis** 714
Thomas A. Stellato, MD; Larry H. Danziger, PharmD; Toni Hau, MD, PhD;
Michael W. L. Gauderer, MD; John L. Ferron, MD; Nahida Gordon, PhD, Cleveland
Single-agent antibiotic therapy with moxalactam was as effective as
tobramycin-clindamycin in treating patients with intra-abdominal infection.
Editorial Comment by H. Harlan Stone, MD, Cleveland

The ARCHIVES OF SURGERY (ISSN 0004-0010) is published monthly by the American Medical Association, 535 N Dearborn St, Chicago, IL 60610, and is an official publication of the Association. Second-class postage paid at Chicago, IL 60610 and at additional mailing office.

SUBSCRIPTION RATES—The subscription rates for the Archives of Surgery are as follows: \$40 for one year, \$70 for two years in the United States and US possessions; all other countries, one year, \$52; two years, \$96 (add \$15 surcharge for each one-year subscription, \$30 for two years for air delivery to the United Kingdom, Europe, Bangladesh, India, Nepal, or Sri Lanka). Rates for subscriptions for delivery to Australia, Japan, New Zealand, or South Korea are available through respective agents. **Special rates for residents and medical students in the United States and US possessions are available.** Address inquiries to the American Medical

Association, Circulation and Fulfillment Division, 535 N Dearborn St, Chicago, IL 60610. Phone: (312) 280-7168.

CHANGE OF ADDRESS—POSTMASTER, send all address changes to Caryl L. Wertheimer, Director of Fulfillment, 535 N Dearborn St, Chicago, IL 60610. Notification of address change must be made at least six weeks in advance, include both old and new addresses, a recent mailing label, and your new zip code.

ADVERTISING PRINCIPLES: Each advertisement in this issue has been reviewed and complies with the principles governing advertising in AMA scientific publications. A copy of these principles is available on request. The appearance of advertising in AMA publications is not an AMA guarantee or endorsement of the product or the claims made for the product by the manufacturer.

CONTENTS—Continued

Hemobilia: A Disease in Evolution	718
Abraham Czerniak, MD; Jeremy N. Thompson, MA, FRCS; Ann P. Hemingway, FRCR; Odd Soreide, MD; Irving S. Benjamin, FRCS; David J. Allison, MD, FRCR; Leslie H. Blumgart, MD, FRCS, London Selective hepatic arterial embolization is the treatment of choice for hemobilia.	
Postoperative Acute Gastrointestinal Tract Hemorrhage and Multiple-Organ Failure	722
Eduardo Bumaschny, MD; Guillermo Doglio, MD; Jorge Pusajó, MD; Liliana Vetere, MD; Carlos Parra, MD; Ricardo M. Grosso, MD; Eduardo Schieppati, MD, Buenos Aires In this report from Argentina, postoperative upper GI tract bleeding was associated with sepsis and multiple-organ failure.	
Computed Tomography of Abdominal Aortic Aneurysm: An In Vivo Pathological Report With a Note on Dynamic Predictors	727
George Pillari, MD; John B. Chang, MD; Joseph Zito, MD; Jon R. Cohen, MD; Kenneth Gersten, MD; Anthony Rizzo, MD; Ariadne M. Bach, MD, New Hyde Park, NY The relationship of thrombus volume and thrombus disposition to rupture remains obscure.	
Marijuana and Alcohol Use Among 1023 Trauma Patients: A Prospective Study	733
Carl A. Soderstrom, MD; Anna L. Trifillis, PhD; Belavadi S. Shankar, DSc; William E. Clark, MS; R Adams Cowley, MD, Baltimore Substance abuse is a pervasive contributor to the occurrence of trauma. Editorial Comment by Donald D. Trunkey, MD, Portland, Ore	
Lower Gastrointestinal Hemorrhage in Renal Transplant Recipients	739
Steven Stylianou, MD; Kenneth A. Forde, MD; Alan I. Benvenisty, MD; Mark A. Hardy, MD, New York Bleeding was caused by pseudomembranous, ischemic, uremic, or opportunistic infectious colitis or idiopathic ulcers. Invited Editorial Comment by Robert J. Corry, MD, Iowa City	
Hand Infections: Bacteriology and Treatment: A Prospective Study	745
E. Patchen Dellinger, MD; Margaret J. Wertz, RN, MN; Stephen D. Miller, MD; Marie B. Coyle, PhD, Seattle Penicillinase-resistant penicillin antibiotics are recommended for hand infections, except for those caused by human bites, which require second-generation cephalosporins. Invited Editorial Comment by Martin C. Robson, MD, Detroit	
Abnormal Rabbit Heterophil Chemotaxis Following Thermal Injury: An In Vivo Model of an Abnormality of the Chemoattractant Receptor for f-met-leu-phe	752
John Mihran Davis, MD, New York, John I. Gallin, MD, Washington, DC Following thermal injury there was decreased granule content and increased chemoattractant binding to rabbit heterophils (neutrophils).	
Balloon Dilatation of Anastomotic Strictures	759
Pat W. Whitworth, MD; Ronald L. Richardson, MD; Gerald M. Larson, MD, Louisville Upper gastrointestinal tract strictures were treated by balloon dilatation.	
Peritoneal Mesothelioma	763
William J. Plaus, MD, Denver Widespread peritoneal neoplastic growth was the common finding at laparotomy.	
CLINICAL NOTES	
Variceal Bleeding, Hypersplenism, and Systemic Mastocytosis: Pathophysiology and Management	767
Bauer E. Sumpio, MD, PhD; Gloria O'Leary, MD; Richard J. Gusberg, MD, New Haven, Conn Mast cell infiltration of the liver or spleen may result in portal hypertension and variceal bleeding.	
Surgical Therapy in Two Patients With Pedunculated Hepatocellular Carcinoma	772
Mark W. Moritz, MD, Detroit; Mamoru Shoji, MD, Atlanta; Gregorio A. Sicard, MD, St Louis; Ryuzo Shioda, MD, Kyoto, Japan; Katherine DeSchryver, MD, St Louis Despite resection in these patients, they died at three and seven months.	
Complications of Needle Localization of Foreign Bodies and Nonpalpable Breast Lesions	775
Andrew D. Bronstein, MD; Ray F. Kilcoyne, MD; Roger E. Moe, MD, Seattle Attempts at needle localization may result in broken and displaced wires.	
BRIEF CLINICAL NOTES	
Obstructive Jaundice: An Unusual Delayed Presentation of Congenital Diaphragmatic Hernia	780
Michael Schiffler, MD; Frederick J. Rescorla, MD; Joseph Fitzgerald, MD; Jay L. Grosfeld, MD, Indianapolis A newborn developed vomiting and jaundice due to an unrecognized right-sided Bochdalek hernia.	
REGULAR DEPARTMENTS	
Correspondence and Brief Communications	782
Books	784
Calendar	787
Instructions for Authors	See May 1988 Issue, p 536
Index to Advertisers	738