

**EDITORIALS**

<b>Hail and Farewell</b> .....	17
Arthur E. Baue, MD, St Louis	

<b>A New Editor for the ARCHIVES</b> .....	18
Michael D. Springer, Chicago	

<b>Inaugural Editorial</b> .....	19
Claude H. Organ, Jr, MD, Oakland, Calif	

**PAPERS READ BEFORE THE ANNUAL MEETING OF THE SOCIETY OF SURGICAL ONCOLOGY, NEW ORLEANS, MAY 22 TO MAY 25, 1988—Part I**

<b>The Prognostic Significance of Lymph Node Metastases After Preoperative Chemotherapy for Locally Advanced Breast Cancer</b> .....	21
--	----

David R. McCready, MD; Gabriel N. Hortobagyi, MD; Shu W. Kau, RN;  
Terry L. Smith, MS; Aman U. Buzdar, MD; Charles M. Balch, MD, Houston

- The number of lymph nodes with metastatic cancer had more prognostic value than other factors.

<b>A Comparison of Palpable and Nonpalpable Breast Cancers</b> .....	26
--	----

Timothy J. Pagana, MD; Willem J. Lubbe, MD; Sheldon M. Schwartz, MD; Gene D. Sprechini, PhD, Williamsport, Pa

- Obviously, nonpalpable breast cancers were smaller and more frequently noninvasive and less frequently had metastasized to lymph nodes.

<b>Nonpalpable In Situ Ductal Carcinoma of the Breast: Predictors of Multicentricity and Microinvasion and Implications for Treatment</b> .....	29
---	----

Gordon F. Schwartz, MD; Arthur S. Patchefsky, MD; Sidney D. Finklestein, MD;  
Sae H. Sohn, MD; Anthony Prestipino, MD; Stephen A. Feig, MD; Jodi S. Singer, MD, Philadelphia

- Ductal carcinoma in situ as an incidental finding may be treated by excision alone.

<b>Breast Carcinoma In Situ</b> .....	33
---------------------------------------	----

David W. Kinne, MD; Jeanne A. Petrek, MD; Michael P. Osborne, MD;  
Aifred A. Fracchia, MD; Angelo A. DePalo, MD; Paul Peter Rosen, MD, New York

- Axillary dissection is not indicated for in situ breast cancer.

<b>Implications of Accurate Pathologic Margins in the Treatment of Primary Breast Cancer</b> .....	37
--	----

Thomas G. Frazier, MD, Bryn Mawr, Pa; Randolph W. Y. Wong, MD, Philadelphia; David Rose, MD, Bryn Mawr, Pa

- Patients with clear or close margins after segmental resection of breast cancer were found to have residual tumor with subsequent reexcision.

<b>Labiality of Steroid Hormone Receptors Following Devascularization of Breast Tumors</b> .....	39
--	----

Lee M. Ellis, MD, Gainesville, Fla; James L. Wittliff, PhD, Louisville; Michael S. Bryant, MD, Harry S. Sitren, PhD, Gainesville, Fla;  
William E. Hogancamp, Louisville; Wiley W. Souba, MD, ScD, Kirby I. Bland, MD, Gainesville, Fla

- Thirty minutes of tissue ischemia before processing resulted in a decrease in estrogen and androgen receptors.

<b>Cancer Centers—Origins and Purpose: The James Ewing Lecture</b> .....	43
--	----

William W. Shingleton, MD, Durham, NC

- Comprehensive cancer centers have been important, but continuous evaluation of their effectiveness is necessary.

<b>Results of Surgical Excision of One to 13 Hepatic Metastases in 98 Consecutive Patients</b> .....	46
--	----

John P. Minton, MD, PhD; Walter B. Hamilton; Armando Sardi, MD; Carol Nieroda, MD;  
Brenda Sickle-Santanello, MD; Patrick J. O'Dwyer, MD, Columbus, Ohio

- Resection of multiple liver metastases by local removal using electrocautery was associated with prolonged survival in a number of patients.

<b>Experimental Liver Metastasis: Implications of Clonal Proclivity and Organ Specificity</b> .....	49
---	----

T. S. Ravikumar, MD; John D'Emilia, MD; Celia Cocchiari; Barbara Wolf, MD; Vincent King; Glenn Steele, Jr, MD, PhD, Boston

- Spontaneous metastases to lung and liver were established in this animal model.

The **ARCHIVES OF SURGERY** (ISSN 0004-0010) is published monthly by the American Medical Association, 535 N Dearborn St, Chicago, IL 60610, and is an official publication of the Association. Second-class postage paid at Chicago, IL 60610 and at additional mailing office.

**SUBSCRIPTION RATES**—The subscription rates for the ARCHIVES OF SURGERY are as follows: \$50 for one year, \$85 for two years in the United States and US possessions; all other countries, one year, \$60; two years, \$105 for surface delivery. (For expedited air delivery to most countries, add \$15 surcharge for one-year subscription, \$30 for two years.) Rates for subscriptions for delivery to Japan or South Korea are available through exclusive agents—contact the publisher. **Special rates for residents and medical students in the United States and US possessions are available.** Address inquiries to the American Medical Association, Circulation and

Fulfillment Division, 535 N Dearborn St, Chicago, IL 60610. Phone: (312)280-7168.

**CHANGE OF ADDRESS**—POSTMASTER, send all address changes for the ARCHIVES OF SURGERY to Caryl L. Wertheimer, Director of Fulfillment, 535 N Dearborn St, Chicago, IL 60610. Notification of address change must be made at least six weeks in advance, include both old and new addresses, a recent mailing label, and your new zip code.

**ADVERTISING PRINCIPLES:** Each advertisement in this issue has been reviewed and complies with the principles governing advertising in AMA scientific publications. A copy of these principles is available on request. The appearance of advertising in AMA publications is not an AMA guarantee or endorsement of the product or the claims made for the product by the manufacturer.

**CONTENTS—Continued**

**Intra-abdominal Recurrence of Colorectal Cancer Detected by Radioimmunoguided Surgery (RIGS System) . . . . . 55**  
Armando Sardi, MD; Mark Workman; Cathy Mojzisek, RN, MS; George Hinkle, RPh; Carol Nieroda, MD;  
Edward W. Martin, Jr, MD, Columbus, Ohio

- In some patients undergoing second-look operations, recurrent tumor was located only by the technique of radiolabeled monoclonal antibodies.

**Rationale for Aggressive Colonoscopy in Patients With Colorectal Neoplasia . . . . . 63**  
Kumar S. Dasmahapatra, MD, Kevin Lopyan, MD, Newark, NJ

- Metachronous colon cancers were found by colonoscopy as late as 60 months after an initial cancer resection.

**Prospective Phase I Trial of Conservative Management of Low Rectal Lesions . . . . . 67**  
David R. McCreedy, MD; David M. Ota, MD; Tyvin A. Rich, MD; Diva Thielvoldt, RN; J. Milburn Jessup, MD, Houston

- In selected patients with low rectal cancer, local excision and radiotherapy was done with early, good results.

**Staging Renal Carcinoma: What Is Sufficient? . . . . . 71**  
Mark A. Benson, MD; John R. Haaga, MD; Martin I. Resnick, MD, Cleveland

- Computed tomography is the principal technique for overall staging of renal cell carcinoma.

**Photodynamic Therapy for Esophageal Tumors . . . . . 74**  
James S. McCaughan, Jr, MD; Thomas A. Nims, MD; Jerry T. Guy, MD; William J. Hicks, MD;  
Thomas E. Williams Jr, MD; Leslie R. Laufman, MD, Columbus, Ohio

- Improvement in ability to swallow was provided in a number of patients.

**Effect of Timing of Lymphokine Presentation on Generation of Cytotoxic T Lymphocytes . . . . . 81**  
Monica M. Stringfellow, MD; Richard E. Wilson, MD; Steven J. Burakoff, MD; Steven H. Herrmann, MD, Boston

- Both interleukin 1 and interleukin 6 are required for optimal generation of cytotoxic T lymphocytes.

**Natural Killer Sensitivity of Colorectal Carcinoma Targets: Correlation With Degree of Differentiation . . . . . 89**  
Darius J. Bāgli, MD, CM; Glenn D. Steele, Jr, MD, PhD; Teresa Barlozzari, PhD, Boston

- Poorly differentiated colorectal cancer cells were susceptible to natural killer cell lysis.

**Cachectin Activity in the Serum of Cachectic, Tumor-Bearing Rats . . . . . 94**  
Mark C. Stovroff, MD; Douglas L. Fraker, MD; Jeffrey A. Norton, MD, Bethesda, Md

- Serum cachectic activity correlated with tumor burden, and the levels became undetectable following tumor resection.

**The Flip Side of Tumor Immunity . . . . . 102**  
Richmond T. Prehn, MD, Liisa M. Prehn, MD, Seattle

- Immunosuppression may be as likely to have a therapeutic effect as immunoaugmentation.

**Recombinant Human Tumor Necrosis Factor Alone and With Chemotherapeutic Agents: Effect on Nude Mouse-Supported Human Bladder Cancer Heterografts . . . . . 107**  
Anurag K. Das, MD; Philip J. Walther, MD, PhD; Niall J. Buckley, MD; Susan H. M. Poulton, Durham, NC

- Synergism between recombinant human tumor necrosis factor and chemotherapeutic agents was found only with etoposide.

**Squamous Cell Carcinoma Arising in Previously Burned or Irradiated Skin . . . . . 115**  
Michael J. Edwards, MD; Rose M. Hirsch, MD; J. Ralph Broadwater, MD; David T. Netscher, MD; Frederick C. Ames, MD, Houston

- Chronic ulceration in scar tissue should suggest this diagnosis.

**Is Limb-Sparing Surgery Applicable to Neurogenic Sarcomas of the Extremities? . . . . . 118**  
John S. Bolton, MD; Jean-Nicolas Vauthey, MD; Gist H. Farr, Jr, MD; Edward I. Sauter, MD;  
John C. Bowen III, MD; David G. Kline, MD, New Orleans

- Patients with high-grade neurogenic sarcomas of the extremities have a high risk of local recurrence after limb-sparing operative treatment.

**Patterns of Recurrence After Resection of Osteosarcoma of the Extremity: Strategies for Treatment of Metastases . . . . . 122**  
James F. Huth, MD, Frederick R. Eilber, MD, Los Angeles

- The prognosis is particularly bad with metastases to the lung and other sites.

**Clinical Experience With Regional Pancreatectomy for Adenocarcinoma of the Pancreas . . . . . 127**  
William F. Sindelar, MD, PhD, Bethesda, Md

- Regional pancreatectomy with en bloc pancreatectomy and removal of regional nodes, duodenum, gastric antrum, and intrapancreatic portal vein produced little survival benefit for most patients.

**REGULAR DEPARTMENTS**

**Correspondence and Brief Communications . . . . . 133**

**Books . . . . . 134**

**Calendar of Events . . . . . 135**

**Instructions for Authors . . . . . 10**

**Index to Advertisers . . . . . 13**