

Copyright © 1989 by the AMERICAN MEDICAL ASSOCIATION

Official Publication of the Western Surgical Association,
Surgical Infection Society, New England Surgical Society,
Society of Surgical Oncology, Pacific Coast Surgical Association**PAPERS READ BEFORE THE ANNUAL MEETING OF THE SOCIETY
OF SURGICAL ONCOLOGY, NEW ORLEANS, LA, MAY 22 TO MAY 25, 1988—Part II**

- Conservative Surgery and Radiation Therapy for Early Breast Cancer: Long-term Cosmetic Results** 153
Mary Ann Rose, MD, Boston, Mass; Ivo Olivotto, MD, Vancouver, Canada; Blake Cady, MD; Clinton Koufman, MD;
Robert Osteen, MD; Barbara Silver; Abram Recht, MD; Jay R. Harris, MD, Boston, Mass
• Satisfactory cosmetic results after conservative operations and radiation therapy are maintained for at least seven years.
- Local Recurrent Mammary Carcinoma Failing Multimodality Therapy: A Solution** 158
Patricia M. McCormack, MD; Manjit S. Bains, MD; Michael E. Burt, MD, PhD;
Nael Martini, MD; Ted Chaglassian, MD; David A. Hidalgo, MD, New York, NY
• Chest wall resection and reconstruction offer significant palliation.
- Extent of Lymph Node Dissection in Melanoma of the Trunk or Lower Extremity** 162
Daniel G. Coit, MD, Murray F. Brennan, MD, New York, NY
• More extensive lymphadenectomy was not associated with improved survival or decreased recurrence.
- Hepatectomy Prolongs Survival of Mice With Induced Liver Metastases** 167
Manuel H. Castillo, MD; Ralph J. Doerr, MD; Norman Paolini, Jr, MS; Stephen Cohen, PhD; Martin Goldrosen, PhD, Buffalo, NY
• Hepatectomy prolonged survival when performed after tumor cell injection.
- Partial Hepatectomy Augments the Liver's Antitumor Response** 170
Ralph Doerr, MD; Manuel Castillo, MD; Paul Evans, MS; Norman Paolini, MS;
Martin Goldrosen, PhD; Stephan A. Cohen, PhD, Buffalo, NY
• Partial hepatectomy enhanced the antitumor activity of nonparenchymal liver cells.
- Prolongation of Survival of Nude Mice Bearing Human Colon Cancer: Treatment With Yttrium 90–Labeled Anti–Carcinoembryonic Antigen Antibody** 175
David M. Hyams, MD; Jose M. Esteban, MD; Barbara G. Beatty, PhD, Duarte, Calif;
Philip M. Wanek, RPh, San Diego, Calif; J. David Beatty, MD, Duarte, Calif
• Different dosing regimens had very different results in this animal model.
- Survival of Patients With Stage B2 Colon Carcinoma: The Gastrointestinal Tumor Study Group Experience** 180
Russell Nauta, MD, Washington, DC; Donald M. Stablein, PhD, Potomac, Md; E. Douglas Holyoke, MD, Buffalo, NY
• Because of excellent survival, patients with stage B2 colon carcinoma are seldom candidates for adjuvant trials of toxic therapy.
- Interactions of Human Colorectal Carcinoma Cells With Basement Membranes: Analysis and Correlation With Differentiation** 183
George W. Daneker, Jr, MD; Andrew J. Piazza; Glenn D. Steele, Jr, MD, PhD; Arthur M. Mercurio, PhD, Boston, Mass
• Tumor cell adherence and poor differentiation correlate with invasion of basement membranes.

The ARCHIVES OF SURGERY (ISSN 0004-0010) is published monthly by the American Medical Association, 535 N Dearborn St, Chicago, IL 60610, and is an official publication of the Association. Second-class postage paid at Chicago, IL 60610 and at additional mailing office.

SUBSCRIPTION RATES—The subscription rates for the ARCHIVES OF SURGERY are as follows: \$50 for one year, \$85 for two years in the United States and US possessions; all other countries, one year, \$60; two years, \$105 for surface delivery. (For expedited air delivery to most countries, add \$15 surcharge for one-year subscription, \$30 for two years.) Rates for subscriptions for delivery to Japan or South Korea are available through exclusive agents—contact the publisher. **Special rates for residents and medical students in the United States and US possessions are available.** Address inquiries to the American Medical Association, Circulation and

Fulfillment Division, 535 N Dearborn St, Chicago, IL 60610. Phone: (312)280-7168.

CHANGE OF ADDRESS—POSTMASTER, send all address changes for the ARCHIVES OF SURGERY to Caryl L. Wertheimer, Director of Fulfillment, 535 N Dearborn St, Chicago, IL 60610. Notification of address change must be made at least six weeks in advance, include both old and new addresses, a recent mailing label, and your new zip code.

ADVERTISING PRINCIPLES: Each advertisement in this issue has been reviewed and complies with the principles governing advertising in AMA scientific publications. A copy of these principles is available on request. The appearance of advertising in AMA publications is not an AMA guarantee or endorsement of the product or the claims made for the product by the manufacturer.

CONTENTS—Continued

The Desmoid Tumor: Not a Benign Disease 191
Mitchell C. Posner, MD; Man H. Shiu, MD; James L. Newsome, MD; Steven I. Hajdu, MD;
Jeffrey J. Gaynor, PhD; Murray F. Brennan, MD, New York, NY
● Desmoid tumors are malignant soft-tissue neoplasms that require aggressive therapy.

Scar or Recurrent Rectal Cancer: Positron Emission Tomography Is More Helpful for Diagnosis Than Immunoscintigraphy 197
Peter Schlag, MD; Burkhard Lehner, MD; Ludwig G. Strauss, MD; Peter Georgi, MD;
Christian Herfarth, MD, Heidelberg, Federal Republic of Germany
● All patients with increased uptake of flu-deoxyglucose F 18 by positron emission tomography had recurrent rectal cancer.

Prognostic Significance of Cellular Immunity to Autologous Breast Carcinoma and Glycoprotein 55 202
Maurice M. Black, MD; Reinhard E. Zachrau, MD; Roy H. Ashikari, MD, Valhalla, NY; Benjamin F. Hankey, ScD, Bethesda, Md
● Cellular immunity as measured by skin window reactivity to autologous breast cancer is prognostically favorable.

Photodynamic Therapy for Cutaneous and Subcutaneous Malignant Neoplasms 211
James S. McCaughan, Jr, MD; Jerry T. Guy, MD; William Hicks, MD;
Leslie Laufman, MD; Thomas A. Nims, MD; John Walker, Columbus, Ohio
● Photodynamic therapy produced a complete response with no evidence of tumor in a number of patients.

Antioxidants vs Carotenoids: Inhibitors or Promoters of Experimental Colorectal Cancers 217
Thomas A. Colacchio, MD; Vincent A. Memoli, MD; Laurie Hildebrandt, Hanover, NH
● Ascorbic acid and alpha-tocopherol demonstrated a chemopreventive effect against induced murine tumors.

Plasma Ultrafiltration as Cancer Therapy 222
Alan Wile, MD, Orange, Calif
● Serum ultrafiltration therapy in six patients reduced their serum suppressor factors.

Endoscopic Ablation of Esophageal Malignancies With the Neodymium-YAG Laser and Electrofulguration 225
Hector R. Nava, MD; Marlys E. Schuh, MD; Raman Nambisan, MD; John L. Clark, MD; Harold O. Douglass, Jr, MD, Buffalo, NY
● Both techniques improved swallowing with mean survival for these patients of 11 months.

Are Complications in Intraoperative Radiation Therapy More Frequent Than in Conventional Treatment? 229
Douglas T. Cromack, MD; Michelle M. Maher, RN; Harald Hoekstra, MD, PhD;
Timothy J. Kinsella, MD; William F. Sindelar, MD, PhD, Bethesda, Md
● No, complications are not more frequent with intraoperative radiation therapy!

The Influence of Natural Killer Cells in Neuroblastoma 235
John V. Reynolds, FRCS(I); Jian Shou, MD; Hoon Choi, MD; Robert Sigal, MD;
Moritz M. Ziegler, MD; John M. Daly, MD, Philadelphia, Pa
● In vivo depletion of natural killer cells enhanced tumor growth and decreased survival in this animal model.

ORIGINAL ARTICLES

Pseudocysts in Chronic Pancreatitis: Surgical Results in 102 Consecutive Patients 240
Tuula Kiviluoto, MD; Leena Kivisaari, MD; Eero Kivilaakso, MD; Matti Lempinen, MD, Helsinki, Finland
● In patients with chronic pancreatitis and pseudocysts, internal drainage or resection produced the most satisfactory results.

Single-Dose Antibiotic Prophylaxis in Contaminated Abdominal Surgery 244
John C. Hall, FRACS, Perth, Western Australia; J. McK. Watts, FRACS, Lexi Press, RN, Bedford Park, South Australia;
P. O'Brien, FRACS, Melbourne, Australia; J. Turnidge, FRACP, P. McDonald, FRACP, Bedford Park, South Australia
● In patients undergoing contaminated abdominal operations, a single dose of prophylactic antibiotic was as effective as an extended course of antibiotics.

Long-term Survival in Carcinoma of the Biliary Tract: Analysis of Prognostic Factors in 146 Resections 248
Kiyooki Ouchi, MD; Seiki Matsuno, MD; Toshio Sato, MD, Sendai, Japan
● Following resection, 41% of patients with gallbladder cancer and approximately 23% of patients with bile duct cancer lived 5 years.

Follow-up of Patients With Aspirated Breast Cysts Is Necessary 253
Hisham Hamed, MD; Andrew Coady, MS; Murid A. Chaudary; Ian S. Fentiman, MD, London, England
● Reexamination six weeks following aspiration of a breast cyst is necessary.

REGULAR DEPARTMENTS

Correspondence and Brief Communications 256

Books 258

Instructions for Authors 148

Index to Advertisers 149