

Copyright © 1989 by the AMERICAN MEDICAL ASSOCIATION

Official Publication of the Western Surgical Association, Surgical Infection Society, New England Surgical Society, Society of Surgical Oncology, Pacific Coast Surgical Association

ORIGINAL ARTICLES

- Evaluation of Prostacyclin Production by Human Gallbladder** 277
 Donald L. Kaminski, MD; Y. G. Deshpande, MS; Scott Westfall, MD; David Herbold, MD, St Louis, Mo
 • Prostacyclin formation may be an important mediator of the inflammatory changes of cholecystitis.
- The Efficacy of Oral Antimicrobials in Reducing Aerobic and Anaerobic Colonic Mucosal Flora** 281
 Jonathan I. Groner, MD; Charles E. Edmiston, Jr, PhD; Candace J. Krepel; Gordon L. Telford, MD;
 Robert E. Condon, MD, MS, Milwaukee, Wis
 • Oral antibiotics with mechanical cleansing, but not mechanical preparation alone, decreased mucosa-associated bacteria.
- Emergency Abdominal Operations in the Patient With Acquired Immunodeficiency Syndrome** 285
 Joshua H. Burack, MD; Marc S. Mandel, MD; Lawrence S. Bizer, MD, Bronx, NY
 • Mortality rate was high following emergency intra-abdominal operations in AIDS patients.
- Errors in Estimating Energy Expenditure in Critically Ill Surgical Patients** 287
 Vincente Cortes, MD, Miami, Fla, Loren D. Nelson, MD, Nashville, Tenn
 • Because of overestimation of the stress factor, calculated or estimated energy expenditure was greater than measured energy expenditure.
- Hemostasis of Solid Viscus Trauma by Intraparenchymal Injection of Fibrin Glue** 291
 Carl J. Hauser, MD, Los Angeles, Calif
 • Intraparenchymal hepatic fibrin glue injection was a more effective hemostatic agent than was topical fibrin glue.
- Neutron Therapy for Nonresectable Radioresistant Tumors** 294
 Lionel Cohen, MD, Frank R. Hendrickson, MD, Batavia, Ill
 • High-energy fast neutron beams are a valuable adjunct to manage some tumors that are resistant to conventional therapy.
- Gallbladder Response to Enteral Lipids in Injured Patients** 301
 Ronald C. Merrell, MD; Priscilla Miller-Crotchett, RN; Patricia Lowry, MD, Houston, Tex
 • After injury these patients had modest enlargement and diminished contractions of the gallbladder that were not improved with enteral lipids.
- Gastric Adenocarcinoma: A Disease in Transition** 303
 Blake Cady, MD, Boston; Ricardo L. Rossi, MD, Mark L. Silverman, MD, Burlington, Mass;
 William Piccione, MD, Chicago, Ill; Thomas A. Heck, MD, Dayton, Ohio
 • Twenty-seven (47%) of 57 patients who underwent resection for cure survived five years.
- Is Extensive Lymphadenectomy Necessary for Surgical Treatment of Intramucosal Carcinoma of the Stomach?** 309
 Keiji Iriyama, MD; Toyomatsu Asakawa, MD; Hiroshi Koike, MD; Hiroshi Nishiwake, MD; Hiroshi Suzuki, MD, Tsu, Japan
 • Lymph node metastases are unlikely in patients with protruding, intramucosal intestinal type gastric carcinomas less than 5 cm in size.
- DNA Ploidy and Tumor Invasion in Human Gastric Cancer: Histopathologic Differentiation** 314
 Daisuke Korenaga, MD; Masaru Haraguchi, MD; Takeshi Okamura, MD; Hideo Baba, MD;
 Atsushi Saito, MD; Keizo Sugimachi, MD, Fukuoka, Japan
 • Aneuploid DNA patterns were associated with more aggressive gastric cancer than were diploid DNA profiles.

The ARCHIVES OF SURGERY (ISSN 0004-0010) is published monthly by the American Medical Association, 535 N Dearborn St, Chicago, IL 60610, and is an official publication of the Association. Second-class postage paid at Chicago, IL 60610 and at additional mailing office.

SUBSCRIPTION RATES—The subscription rates for the ARCHIVES OF SURGERY are as follows: \$50 for one year, \$85 for two years in the United States and US possessions; all other countries, one year, \$60; two years, \$105 for surface delivery. (For expedited air delivery to most countries, add \$15 surcharge for one-year subscription, \$30 for two years.) Rates for subscriptions for delivery to Japan or South Korea are available through exclusive agents—contact the publisher. **Special rates for residents and medical students in the United States and US possessions are available.** Address inquiries to the American Medical Association, Circulation and

Fulfillment Division, 535 N Dearborn St, Chicago, IL 60610. Phone: (312)280-7168.

CHANGE OF ADDRESS—POSTMASTER, send all address changes for the ARCHIVES OF SURGERY to Subscription Department, 535 N Dearborn St, Chicago, IL 60610. Notification of address change must be made at least six weeks in advance, include both old and new addresses, a recent mailing label, and your new zip code.

ADVERTISING PRINCIPLES: Each advertisement in this issue has been reviewed and complies with the principles governing advertising in AMA scientific publications. A copy of these principles is available on request. The appearance of advertising in AMA publications is not an AMA guarantee or endorsement of the product or the claims made for the product by the manufacturer.

CONTENTS—Continued

Prophylactic Antibiotherapy in Abdominal Surgery: First- vs Third-Generation Cephalosporins	323
Nelly Rotman, MD, Creteil, France; Jean-Marie Hay, MD, Colombes, France; Françoise Lacaine, MD, Paris, France; Pierre-Louis Fagniez, MD, Creteil, France; Association de Recherche en Chirurgie Cooperative Group, Bois-Colombes, France	
● Cephalosporin prophylaxis was helpful in clean operations, in clean contaminated operations, and in patients with general increased risk of infection.	
Guidelines for Practical Utilization of Intraoperative Frozen Sections	331
Marianne U. Prey, MD, St Louis, Mo; Thomas Vitale, MD, Melbourne, Fla; Scott A. Martin, MD, St Louis, Mo	
● Intraoperative frozen-section interpretation was associated with 0.15% false-positive and 2.8% false-negative diagnosis rates.	
Comparison of Antibody Response With Delayed Hypersensitivity in Severely Injured Patients	339
Michael J. Hershman, FRCS; William G. Cheadle, MD; Sarah H. Appel; Christopher D. George, FRCS; Pattie F. Davidson, RN; Karen M. Cost, PhD; Hiram C. Polk, Jr, MD, Louisville, Ky	
● Clinical outcome and antibody response were determined by the extent of initial bacterial contamination.	
Computed Tomography and Diagnostic Peritoneal Lavage in Blunt Abdominal Trauma: Their Combined Role	344
Paul A. Kearney, Jr, MD; Thomas Vahey, MD; Richard E. Burney, MD; Gary Glazer, MD, Ann Arbor, Mich	
● Peritoneal lavage was helpful in detecting injuries missed by computed tomography.	
The Training and Utilization of Surgical Physician Assistants: A Retrospective Study	348
Linda Ballard Brandt; Malcolm S. Beinfeld, MD; Horace A. Laffaye, MD, Norwalk, Conn; Arthur E. Baue, MD, St Louis, Mo	
● A surgical postgraduate training program for physician assistants provided coverage of surgical patients in hospitals without surgical residencies.	
Physiologic Effects of Steep Positioning in the Surgical Intensive Care Unit	352
Loren D. Nelson, MD, Hans B. Anderson, RRT, Miami, Fla	
● Continuous rotation reversed the adverse effects on oxygenation caused by positional changes.	
Nosocomial Abscess: Results of an Eight-Year Prospective Study of 32 284 Operations	356
Mary M. Olson, RN, BSN, Melody O'Connor Allen, MD, Minneapolis, Minn	
● Postoperative infection surveillance was associated with a decrease in wound infection and abscess formation rates.	
Determinants of Urea Nitrogen Production in Sepsis: Muscle Catabolism, Total Parenteral Nutrition, and Hepatic Clearance of Amino Acids	362
Mauro Pittiruti, MD; John H. Siegel, MD; Gabriele Sganga, MD; Bill Coleman; Charles E. Wiles III, MD; Robert Placko, Baltimore, Md	
● Septic patients who survived had strikingly different metabolic alterations from septic patients with multiple organ failure.	
Isolated Regional Perfusion in the Treatment of Subungual Melanoma	373
Peter C. Baas, MD; Harald J. Hoekstra, MD, PhD; Heimen Schraffordt Kooops, MD, PhD; J. Wolter Oosterhuis, MD, PhD; Jan Oldhoff, MD, PhD, Groningen, the Netherlands	
● Regional hyperthermic cytotoxicity drug perfusion prevented local recurrence following amputation for subungual melanoma.	
CLINICAL NOTES	
Influence of Estrogen Receptors on Survival and Recurrence in Patients With Breast Cancer Without Lymph Node Metastases	377
Hope S. Maki, MD, James L. Hoehn, MD, Marshfield, Wis	
● Estrogen receptor status did not adversely affect outcome in lymph node-negative breast cancer patients.	
A Foregut Carcinoid Tumor Causing Zollinger-Ellison Syndrome	381
Gordon B. Werbel, MD; Suzanne P. Nelson, MD; Philip G. Robinson, MD; John Anastasi, MD; Raymond J. Joehl, MD; Robert V. Rege, MD, Chicago, Ill	
● A benign gastric carcinoid tumor contained gastrin and produced severe peptic ulcer disease.	
TECHNIQUE	
Fibrin Glue Achieves Hemostasis in Patients With Coagulation Disorders	385
Harry B. Kram, MD; Ragu C. Nathan, MD; Frederick J. Stafford, MD; Arthur W. Fleming, MD; William C. Shoemaker, MD, Los Angeles, Calif	
● Fibrin glue is especially useful in patients with parenchymal organ hemorrhage and disordered coagulation.	
REGULAR DEPARTMENTS	
Correspondence and Brief Communications	388
Books	390
Calendar of Events	391
Index to Advertisers	270
Instructions for Authors	See February Issue, p 148