

Copyright © 1989 by the AMERICAN MEDICAL ASSOCIATION

Official Publication of the Western Surgical Association,
Surgical Infection Society, New England Surgical Society,
Society of Surgical Oncology, Pacific Coast Surgical Association**EDITORIALS**

- W. Dean Warren, MD: October 28, 1924–May 10, 1989** 767
Robert Zeppa, MD, Miami, Fla
- Ben Eiseman, MD** 768
Claude H. Organ, Jr, MD, Oakland, Calif
- The Surgeon and Biliary Lithotripsy** 769
David L. Nahrwold, MD, Chicago, Ill
- Western Surgical Association** 770
Claude H. Organ, Jr, MD, Oakland, Calif
- PAPERS READ BEFORE THE 96TH ANNUAL MEETING OF THE WESTERN SURGICAL ASSOCIATION, SAN DIEGO, CALIF, NOV 14-16, 1988**
- They Call It Competition: Presidential Address** 771
Melvin A. Block, MD, PhD, La Jolla, Calif
- An Analysis of the Reduced Morbidity and Mortality Rates After Pancreaticoduodenectomy** 778
Carlos A. Pellegrini, MD; Christopher F. Heck, MD; Steven Raper, MD; Lawrence W. Way, MD, San Francisco, Calif
- Computed tomographic scan preoperative assessment combined with improved operative technique has lowered the mortality rate associated with the Whipple procedure.
- Percutaneous Cholecystostomy in the Diagnosis and Treatment of Acute Cholecystitis in the High-Risk Patient** 782
Gordon B. Werbel, MD; David L. Nahrwold, MD; Raymond J. Joehl, MD; Robert L. Vogelzang, MD; Robert V. Rege, MD, Chicago, Ill
- Percutaneous cholecystostomy and bile cultures in critically ill patients can avoid the excessive risk of cholecystectomy.
- Endoscopic Cholangiography and Stone Removal Prior to Cholecystectomy: A More Cost-effective Approach Than Operative Duct Exploration?** 787
Greg Van Stiegmann, MD; Nathan W. Pearlman, MD; John S. Goff, MD; John H. Sun, MD; Lawrence W. Norton, MD, Denver, Colo
- Patients requiring cholecystectomy with suspected choledocholithiasis may be treated more cost effectively by ERCP with common duct stone removal prior to cholecystectomy than by cholecystectomy and common duct exploration.
- Less Is Better: The Diagnostic Workup of the Patient With Obstructive Jaundice** 791
Richard Olen, MD; Jack Pickleman, MD; Robert J. Freeark, MD, Maywood, Ill
- If an abdominal ultrasound study in the typical patient with obstructive jaundice discloses distal bile duct obstruction, no further tests are necessary.
- Function of Pancreas Allografts More Than 1 Year Following Transplantation** 796
Francis H. Wright, MD; John L. Smith, MD; Scott A. Ames, MD; Barbara Schanbacher, RN; Robert J. Corry, MD, Iowa City, Iowa
- Pancreatic transplantation achieves a euglycemic state in type I diabetes mellitus patients but warrants a long-term follow-up.
- Changes in Tibial Venous Blood Flow in the Evolving Compartment Syndrome** 801
William G. Jones II, MD; Malcolm O. Perry, MD; Harry L. Bush, Jr, MD, New York, NY
- Interference with tibial venous blood flow is an early event with increased muscle compartment pressures and can be followed up with Doppler venous flow evaluation.
- Myocardial Contusion: When Can the Diagnosis Be Eliminated?** 805
Frank B. Miller, MD; Charles R. Shumate, MD; J. David Richardson, MD, Louisville, Ky
- The electrocardiogram and the clinical course were the primary determinants of the need for cardiac monitoring and therapeutic intervention.
- Intra-abdominal Injury Following Blunt Trauma: Identifying the High-Risk Patient Using Objective Risk Factors** 809
Robert C. Mackersie, MD; Anurag D. Tiwary, MD; Steven R. Shackford, MD; David B. Hoyt, MD, San Diego, Calif
- Of 11 clinical indicators assessed, arterial base deficit, major chest injury, hypotension, and pelvic fractures significantly increased the chance of intra-abdominal injury.

The ARCHIVES OF SURGERY (ISSN 0004-0010) is published monthly by the American Medical Association, 535 N Dearborn St, Chicago, IL 60610, and is an official publication of the Association. Second-class postage paid at Chicago, IL 60610, and at additional mailing office.

SUBSCRIPTION RATES—The subscription rates for the ARCHIVES OF SURGERY are as follows: \$50 for one year, \$85 for two years in the United States and US possessions; all other countries, one year, \$60; two years, \$105 for surface delivery. (For expedited air delivery to most countries, add \$15 surcharge for one year subscription, \$30 for two years.) Rates for subscriptions for delivery to Japan or South Korea are available through exclusive agents—contact the publisher. **Special rates for residents and medical students in the United States and US Possessions are available.** Address inquiries to the American Medical Association, Circulation

and Fulfillment Division, 535 N Dearborn St, Chicago, IL 60610. Phone: (312)280-7168.

CHANGE OF ADDRESS—POSTMASTER, send all address changes to the ARCHIVES OF SURGERY, Attention: Subscription Department, 535 N Dearborn St, Chicago, IL 60610. Notification of address change must be made at least six weeks in advance, include both old and new addresses, a recent mailing label, and your new zip code.

Advertising Principles: Each advertisement in this issue has been reviewed and complies with the principles governing advertising in AMA scientific publications. A copy of these principles is available on request. The appearance of advertising in AMA publications is not an AMA guarantee or endorsement of the product or the claims made for the product by the manufacturer.

CONTENTS—Continued

Fine-Needle Aspiration for Breast Mass Diagnosis	814
William H. Wolberg, MD; Martin A. Tanner, PhD; Wei-Yin Loh, PhD, Madison, Wis	
● A statistically based algorithm for fine-needle biopsy of breast masses reveals clinical parameters with a sensitivity of 0.98, a specificity of 0.94 and a positive predictability of 0.92	
Oncological Aspects of Immediate Breast Reconstruction Following Mastectomy for Malignancy	819
Craig H. Johnson, MD; Jon A. van Heerden, MB, FRCS; John H. Donohue, MD, Rochester, Minn; J. Kirk Martin, Jr, MD, Jacksonville, Fla; Ian T. Jackson, MD, FRCS(Glasg), FRCS(Edin), Duane M. Ilstrup, MS, Rochester, Minn	
● Immediate breast reconstruction should be offered women with operable breast cancer because of the positive psychosocial benefit without a negative impact on patient outcome	
The Outcome Index: A Method of Quality Assurance in the Special Care Area	825
Arthur S. McFee, MD, James Gilbert, MD, San Antonio, Tex	
● A system outcome score (SOS) is suggested as a method of quality assurance for the surgical intensive care unit.	
Nonoperative Treatment of Perforated Duodenal Ulcer	830
Thomas V. Berne, MD, Arthur J. Donovan, MD, Los Angeles, Calif	
● The perforated duodenal ulcer can be treated nonoperatively when a gastroduodenogram documents self-sealing.	
Catheter Aspiration for Simple Pneumothorax: Experience with 114 Patients	833
Ralph E. Delius, MD; Farouk N. Obeid, MD; H. Mathilda Horst, MD; Victor J. Sorensen, MD; John J. Fath, MD; Brack A. Bivins, MD, Detroit, Mich	
● Catheter aspiration is a safe, successful, and cost-effective technique for managing the simple pneumothorax.	
Changing Patterns of Treatment and Survival in Neonates With Meconium Ileus	837
Frederick J. Rescorla, MD; Jay L. Grosfeld, MD; Karen J. West, MD; Dennis W. Vane, MD, Indianapolis, Ind	
● Nonoperative technique can relieve obstruction in uncomplicated meconium ileus and avoid enterostomy in most cases.	
Medullary Carcinoma in Children: Results of Early Detection and Surgery	841
Robert L. Telander, MD; Donald Zimmerman, MD; Glen W. Sizemore, MD; Jon A. van Heerden, MB, ChB; Clive S. Grant, MD, Rochester, Minn	
● All children with multiple endocrine neoplasia type IIb (MEN-IIb) should be studied shortly after birth, and those with MEN-IIa should be studied by 1 year of age.	
ORIGINAL ARTICLES	
Evaluation Of Staging Workup in Malignant Melanoma	847
Tawfiq Khansur, MD; Jane Sanders, MD; Suman K. Das, MD, Jackson, Miss	
● There is no need for routine work-up for metastases in malignant melanoma unless the serum LDH is elevated	
Early Detection of Shock in Critically Ill Patients by Skeletal Muscle Po₂ Assessment	853
Gerald I. J. M. Beerthuizen, MD; R. Jan A. Goris, MD; Ferdinand J. A. Kreuzer, PhD, Nijmegen, the Netherlands	
● Skeletal muscle Po ₂ helped predict the development of shock if inotropic agents had not been given.	
Oxidative Activity of Polymorphonuclear Leukocytes After Thermal Injury	856
Marek K. Dobke, MD, PhD; Edwin A. Deitch, MD; Timothy J. Harnar, MD; Charles R. Baxter, MD, Lubbock, Tex	
● Neutrophils from burn patients have increased resting and decreased zymosan-stimulated oxygen consumption when compared with neutrophils of control patients.	
Posttraumatic Autotransplantation of Spleen Tissue	863
Solly Mizrahi, MD; Amitai Bickel, MD; Mahmud Haj, MD; Ida Lunski, MD; Benjamin Shtamler, MD, Nahariya, Israel	
● Following splenectomy for trauma, spleen tissue was successfully autotransplanted in patients	
Recurrent Stenosis by Duplex Scan Following Carotid Endarterectomy	866
Vincent J. Bertin, MD; Fred R. Plecha, MD; Gregory J. Rogers, MD; Karen MacPherson, RNC; Judith Lund, LPN, Cleveland, Ohio	
● Following carotid endarterectomy a higher incidence of recurrent stenosis was found in female patients.	
CLINICAL NOTES	
Primary Aortoduodenal Fistula: Diagnosis by Computed Tomography	870
Ibrahim M. Ibrahim, MD, Joseph S. Raccuia, MD, Englewood, NJ; Joseph Micale, MD, Asham Zafar, MD, North Bergen, NJ	
● A computed tomographic scan aided in the preoperative diagnosis of abdominal aortic aneurysm with duodenal fistula.	
Hernia Uterus Inguinale Associated With Unicornuate Uterus	872
David C. Elliott, MD; Thomas E. Beam, MD; Thomas S. Denapoli, MD, Washington, DC	
● An incarcerated inguinal hernia contained an ovary, a fallopian tube, and a rudimentary uterine horn.	
TECHNIQUE	
A New Surgical Localization Technique for Biopsy in Patients With Nipple Discharge	874
Arindam Choudhury, MD, Paul A. Wengert, Jr, MD, Harrisburg, Pa; J. Stanley Smith, Jr, MD, Hershey, Pa	
● Dye injection localization was used during operative treatment of nipple discharge.	
REGULAR DEPARTMENTS	
Correspondence and Brief Communications	876
Calendar of Events	876
Index to Advertisers	862
Instructions for Authors	845