

Copyright © 1990 by the AMERICAN MEDICAL ASSOCIATION

Official Publication of the Association of Veterans Administration Surgeons, New England Surgical Society, Pacific Coast Surgical Association, Society of Surgical Oncology, Surgical Infection Society, Western Surgical Association.

## EDITORIAL

- Nobel Laureate: Joseph E. Murray, Clinical Surgeon, Scientist, Teacher** . . . . . 1423  
Maurice J. Jurkiewicz, MD

## SPECIAL ARTICLE

- On the Supply of Physicians** . . . . . 1425  
Peter I. Buerhaus, PhD, RN, George D. Zuidema, MD, Ann Arbor, Mich

## ORIGINAL ARTICLES

- Heart Transplantation in Jehovah's Witnesses: An Initial Experience and Follow-up** . . . . . 1430  
Clay M. Burnett, MD; J. Michael Duncan, MD; J. David Vega, MD;  
James L. Lonquist, MD; Michael S. Sweeney, MD; O. H. Frazier, MD, Houston, Tex

**Invited Commentary:** William A. Gay, Jr, MD, Salt Lake City, Utah

- Cardiac transplants may be safely offered to Jehovah's Witnesses without fear of uniformly poor outcome.

- Ambulatory Cholecystectomy Without Disability** . . . . . 1434  
Walter P. Ledet, Jr, MD, Sulphur, La

- Ambulatory cholecystectomy for properly selected patients is feasible, safe, and cost-effective.

**Invited Commentary:** Ronald P. Fischer, MD, PhD, Houston, Tex

- Postoperative Complications in Patients With Disabling Psychiatric Illnesses or Intellectual Handicaps: A Case-Controlled, Retrospective Analysis** . . . . . 1436  
Bruce S. Cutler, MD, Mitchell P. Fink, MD, Worcester, Mass

- A strategy for the postoperative management is presented for institutionalized patients with emphasis on the prevention of pulmonary complications.

- Needle-Localized Mammographic Lesions: Results and Evolving Treatment Strategy** . . . . . 1441  
H. Richard Alexander, MD; Frank C. Candela, MD; D. David Dershaw, MD; David W. Kinne, MD, New York, NY

- The rate of malignant diagnoses after NLCC was 24% in asymptomatic women and 43% in the contralateral breast after treatment for breast cancer.

- Aerobic and Anaerobic Bacteriology of Wounds and Cutaneous Abscesses** . . . . . 1445  
Itzhak Brook, MD, MSc, Edith H. Frazier, MSc, Bethesda, Md

- The highest rates of anaerobes for wound infections were in the inguinal, buttock, and trunk areas; for abscesses, the rates were highest in the perirectal, external genitalia, neck, and inguinal areas.

- Reinforcement of Colonic Anastomoses With a Laser and Dye-Enhanced Fibrinogen** . . . . . 1452  
Nader Moazami; Mehmet C. Oz, MD; Lawrence S. Bass, MD; Michael R. Treat, MD, New York, NY

- Laser-fibrination reinforcement significantly enhances the early strength of sutured colonic anastomoses.

- Hepatic Vein Reconstruction for Preserving Remnant Liver Function** . . . . . 1455  
Satoshi Nakamura, MD; Shukichi Sakaguchi, MD; Tadashi Kitazawa, MD;  
Shouhachi Suzuki, MD; Kenichi Koyano, MD; Hiroyuki Muro, MD, Hamamatsu, Japan

- Hepatic vein reconstruction allows for extended liver resection when malignancies infiltrate major hepatic veins.

The ARCHIVES OF SURGERY (ISSN 0004-0010) is published monthly by the American Medical Association, 515 N State St, Chicago, IL 60610, and is an official publication of the Association. Second-class postage paid at Chicago, IL 60610, and at additional mailing office.

**SUBSCRIPTION RATES**—The subscription rates for the ARCHIVES OF SURGERY are as follows: \$56 for one year, \$97 for two years in the United States and US possessions; all other countries, one year, \$66; two years, \$117 for surface delivery. For expedited air delivery to most countries, add \$15 surcharge for one year subscription, \$30 for two years. (Rates for subscriptions for delivery to Japan or South Korea are available through exclusive agents—contact the publisher.) **Special rates for residents and medical students in the United States and US possessions are available.** Address inquiries to the American Medical Association, Circulation

and Fulfillment Division, 515 N State St, Chicago, IL 60610. Phone: (312)464-0183.

**CHANGE OF ADDRESS**—POSTMASTER, send all address changes to the ARCHIVES OF SURGERY, Attention: Subscription Department, 515 N State St, Chicago, IL 60610. Notification of address change must be made at least six weeks in advance, include both old and new addresses, a recent mailing label, and your new zip code.

**Advertising Principles:** Each advertisement in this issue has been reviewed and complies with the principles governing advertising in AMA scientific publications. A copy of these principles is available on request. The appearance of advertising in AMA publications is not an AMA guarantee or endorsement of the product or the claims made for the product by the manufacturer.

**CONTENTS—Continued**

**Evaluation of Methods for Detecting Venous Reflux:  
Perspectives in Venous Insufficiency** . . . . . **1463**  
Seshadri Raju, MD, Ruth Fredericks, MD, Jackson, Miss  
● Horizontal techniques of descending venography and nucleotide descending venography have unacceptable low sensitivity and were abandoned.

**Does Survival Depend on the Amount of Autotransplanted Splenic Tissue?** . . . . . **1472**  
M. H. Shokouh-Amiri, MD; S. Rahimi-Saber, MD; C. Palnaes Hansen, MD;  
P. Skov Olsen, MD, PhD, S. Lindkaedr Jensen, MD, PhD, Copenhagen, Denmark  
● Forty percent of the spleen should be autotransplanted to protect optimally against bacterial infection.

**Primary Malignant Peritoneal Mesothelioma:  
A Report of Seven Cases and a Review of the Literature** . . . . . **1477**  
Juan A. Asensio, MD, Philadelphia, Pa; Peter Goldblatt, MD, Neil R. Thomford, MD, Toledo, Ohio  
● Long-term survival may be achieved with correct identification and aggressive management; prevention is linked to avoidance of asbestos exposure.

**Aortic Occlusion and Vascular Isolation Allowing Avascular Hepatic Resection** . . . . . **1482**  
Michael S. Stephen, FRACS; A. G. Ross Sheil, MS, FRCS, FRACS; John F. Thompson, FRACS;  
Timothy Wilson, FRCS, FRCS(Edin), FRACS; Stuart L. Boland, FRCS, FRACS, Sydney, Australia  
● Supraceliac abdominal aortic occlusion and hepatic vascular isolation increases ease of operative management, reduced blood loss, and requires less operating time.

**Management of Perineal Wounds Following Abdominoperineal Resection With Inferior Gluteal Flaps** . . . . . **1486**  
Wilbur L. Baird, MD; T. Roderick Hester, MD; Foad Nahai, MD; John Bostwick III, MD, Atlanta, Ga  
● Using an inferior gluteal myocutaneous flap, 15 of 16 wounds achieved healing following abdominoperineal resection for inflammatory bowel disease.

**BASIC SCIENCE REVIEW**

**Antibiotic Pharmacokinetics in Surgery** . . . . . **1490**  
Donald E. Fry, MD, David E. Pitcher, MD, Albuquerque, NM

**SURGICAL HISTORY**

**A History of the Bilioenteric Anastomosis** . . . . . **1493**  
Steven A. Ahrendt, MD, Henry A. Pitt, MD, Baltimore, Md

**CLINICAL NOTES**

**The Papillary-Cystic Neoplasm of the Pancreas:  
An Increasingly Recognized Clinicopathologic Entity** . . . . . **1502**  
Juan A. Sanchez, MD; Kurt D. Newman, MD;  
Martin R. Eichelberger, MD; Russell J. Nauta, MD, Washington, DC  
● Papillary cystic neoplasms of the pancreas occur predominantly in women, have a low malignant potential, and have a high curable rate with resection.

**Extrahepatic Biliary Obstruction by Lymphoma** . . . . . **1507**  
Edward Feller, MD, Fred J. Schiffman, MD, Providence, RI  
● Two patients with obstructive jaundice were satisfactorily palliated without operation when ERCP identified lymphoma as the cause.

**REGULAR DEPARTMENTS**

**Correspondence and Brief Communications** . . . . . **1516**

**Books** . . . . . **1516**

**Instructions for Authors** . . . . . **1519**

**Index to Advertisers** . . . . . **1520**

**Classified Advertising** . . . . . **1521**