

EDITORIAL

- The Future of General Surgery** 145
Claude H. Organ, Jr, MD, Oakland, Calif

COMMENTARIES

- The Surgical Residency: The Job Description Does Not Fit the Job** 147
Walter J. Pories, MD, Hazel M. Aslakson, EdD, Greenville, NC
- The Family History and Cancer Control: Hereditary Breast Cancer** 151
Henry T. Lynch, MD, Omaha, Neb

PAPERS READ BEFORE THE ANNUAL MEETING OF THE SOCIETY OF SURGICAL ONCOLOGY, SAN FRANCISCO, CALIF, MAY 21 TO MAY 24, 1989—PART I

- The Society of Surgical Oncology at a Crossroads: Thoughts for the Future: Presidential Address** 153
Blake Cady, MD, Boston, Mass
- Whither the Solid Tumors: James Ewing Lecture** 158
W. P. Longmire, Jr, MD, Los Angeles, Calif
- Strategies Toward the Primary Prevention of Cancer: Lucy Wortham James Clinical Research Award Lecture** 163
Ernst L. Wynder, MD, New York, NY
- Education in schools and health promotion to all community segments are primary preventive strategies for cancer control.
- Breast Biopsy for Calcifications in Nonpalpable Breast Lesions: A Prospective Study** 170
Dido Franceschi, MD; Joseph Crowe, MD; Robert Zollinger, MD; Renate Duchesneau, MD; Robert Shenk, MD; Gladys Stefanek, RN; Jerry M. Shuck, MD, DSc, Cleveland, Ohio
- A marked correlation is observed between malignancy and small lesions, more than 15 calcifications, and calcifications in a linear or branching pattern.
- The Use of Fine-Needle Aspirates of Breast Cancers to Evaluate Hormone-Receptor Status** 174
Joel Lundy, MD, Stony Brook, NY; Mary Lozowski, MS; Darya Sadri, MS; Yousri Mishriki, MD, Mineola, NY
- Immunocytochemical assay can distinguish receptor-negative and receptor-positive tumors and can be utilized to assay tumors too small for standard steroid analysis.
- Improved Detection of Metastases to Lymph Nodes and Estrogen Receptor Determination** 179
Shepherd H. Schurman; Nehama Sharon, PhD; Robert A. Goldschmidt, MD; Edward F. Scanlon, MD, Evanston, Ill
- A new method of detecting nodal micrometastases in breast cancer combines the advantages of serial sectioning and immunohistochemical staining.
- Tumor-Associated Antigen Immune Complexes: A Potential Marker of Recurrent Melanoma** 187
Jan H. Wong, MD; Shi-Hong Xu; Rishab K. Gupta, PhD; Donald L. Morton, MD, Los Angeles, Calif
- Monitoring tumor-associated antigen-specific immune complexes appears to be a potentially useful marker in human melanoma.
- Patterns of Human Tumor-Infiltrating Lymphocytes in 120 Human Cancers** 200
Charles M. Balch, MD; Lee B. Riley, MD; Yoon Joo Bae, MD; Marie A. Salmeron, MD; Chris D. Platsoucas, PhD; Andrew von Eschenbach, MD; Kyogo Itoh, MD; Houston, Tex
- Tumor-infiltrating lymphocytes from 120 human cancers show considerable diversity with respect to lymphocyte infiltration, phenotypic expression, and functional capacity.
- Metastatic Potential of Colon Carcinoma: Expression of ABO/Lewis-Related Antigens** 206
Stuart D. Hoff, MD; Tatsuro Irimura, PhD; Yoshifumi Matsushita, MD; David M. Ota, MD; Karen R. Cleary, MD, Houston, Tex; Sen-itiroh Hakomori, MD, Seattle, Wash
- Expression of sialyl-dimeric Le^x increases as the metastatic potential of colorectal carcinoma increases and is related to its progression.
- Occult Breast Cancer Presenting With Axillary Metastases: Updated Management** 210
Paul L. Baron, MD; Michael P. Moore, MD, PhD; David W. Kinne, MD; Frank C. Candela, MD; Michael P. Osborne, MD, MS; Jeanne A. Petrek, MD, New York, NY
- Both breast conservation and adjuvant therapy have a role in the management of occult stage II breast cancer.

The ARCHIVES OF SURGERY (ISSN 0004-0010) is published monthly by the American Medical Association, 535 N Dearborn St, Chicago, IL 60610, and is an official publication of the Association. Second-class postage paid at Chicago, IL 60610, and at additional mailing office.

SUBSCRIPTION RATES—The subscription rates for the ARCHIVES OF SURGERY are as follows: \$56 for one year, \$97 for two years in the United States and US possessions; all other countries, one year, \$66; two years, \$117 for surface delivery. For expedited air delivery to most countries, add \$15 surcharge for one year subscription, \$30 for two years. (Rates for subscriptions for delivery to Japan or South Korea are available through exclusive agents—contact the publisher.) **Special rates for residents and medical students in the United States and US Possessions are available.** Address inquiries to the American Medical Association, Circulation

and Fulfillment Division, 535 N Dearborn St, Chicago, IL 60610. Phone: (312)280-7168.

CHANGE OF ADDRESS—POSTMASTER, send all address changes to the ARCHIVES OF SURGERY, Attention: Subscription Department, 535 N Dearborn St, Chicago, IL 60610. Notification of address change must be made at least six weeks in advance, include both old and new addresses, a recent mailing label, and your new zip code.

Advertising Principles: Each advertisement in this issue has been reviewed and complies with the principles governing advertising in AMA scientific publications. A copy of these principles is available on request. The appearance of advertising in AMA publications is not an AMA guarantee or endorsement of the product or the claims made for the product by the manufacturer.

CONTENTS—Continued

| | |
|---|------------|
| Changes Associated With Metastasis in B16-F1 Melanoma Cells Surviving Heat | 216 |
| S. David Nathanson, MD; Robert F. Cerra, PhD; Fred W. Hetzel, PhD; Richard J. Zarbo, MD; John D. Crissman, MD; Robert Page, MS; Patricia Anaya, MS; Patricia Westrick, Detroit, Mich | |
| ● After subcurative hyperthermia, increased expression of putative laminin receptors may be associated with increased metastasis of melanomas. | |
| Successful Combination Immunotherapy for the Generation In Vivo of Antitumor Activity With Anti-CD3, Interleukin 2, and Tumor Necrosis Factor | 220 |
| Stephen C. Yang, MD; Kim D. Fry; Elizabeth A. Grimm, PhD; Jack A. Roth, MD, Houston, Tex | |
| ● The sequential use of anti-CD3, interleukin 2, and tumor necrosis factor for lymphokine-activated killer cells induction and maintenance potentiates antitumor activity. | |
| Indium 111 ZCE-025 Immunoscintigraphy in Occult Recurrent Colorectal Cancer With Elevated Carcinoembryonic Antigen Level | 226 |
| Ralph J. Doerr, MD, Hani Abdel-Nabi, MD, PhD, Buffalo, NY; Bruce Merchant, MD, PhD, San Diego, Calif | |
| ● Indium 111 labeled carcinoembryonic antigen monoclonal antibody scanning is a useful diagnostic localizing agent for recurrent colorectal metastases. | |
| Preoperative Staging and Assessment of Resectability of Pancreatic Cancer | 230 |
| Andrew L. Warshaw, MD; Zhuo-yun Gu, MD; Jack Wittenberg, MD; Arthur C. Waltman, MD, Boston, Mass | |
| ● Accurate and efficient preoperative staging of pancreatic and periampullary adenocarcinoma with existing diagnostic modalities can be accomplished. | |
| Carcinoma of the Gallbladder: Does Radical Resection Improve Outcome? | 237 |
| John H. Donohue, MD; David M. Nagorney, MD; Clive S. Grant, MD; Kenichi Tsushima, MD; Duane M. Ilstrup, MS; Martin A. Adson, MD, Rochester, Minn | |
| ● Though radical cholecystectomy may benefit individual patients, there was no survival advantage when compared with cholecystectomy. | |
| A Phase I Trial of Preoperative Radiation, Proctectomy, and Endoanal Reconstruction | 247 |
| Alfred M. Cohen, MD, Bruce D. Minsky, MD, New York, NY | |
| ● Phase I studies are reported on candidates for abdominoperineal resection combined with preoperative radiation therapy, restorative proctectomy, and endoanal anastomosis. | |
| Invited Commentary: Malcolm C. Veidenheimer, MD, Burlington, Mass | |
| Surgical Implications of Primary Gastrointestinal Lymphoma of Childhood | 252 |
| Irvin D. Fleming, MD; Peter S. Turk, MD; Sharon B. Murphy, MD; William M. Crist, MD; Victor M. Santana, MD; Bhaskar N. Rao, MD, Memphis, Tenn | |
| ● When feasible, children with primary non-Hodgkin's lymphoma of the gastrointestinal tract benefit from complete resection. | |
| Invited Commentary: James A. O'Neill, Jr, MD, Philadelphia, Pa | |
| The Genetic Bases of Cancer: Lucy Wortham James Lecture | 257 |
| Robert A. Weinberg, PhD, Cambridge, Mass | |
| ORIGINAL ARTICLES | |
| Predictive Factors for Bactibilia in Acute Cholecystitis | 261 |
| Jesse E. Thompson, Jr, MD, Robert S. Bennion, MD, Sylmar, Calif; Jeffery E. Doty, MD, Elvira L. Muller, MD, Sepulveda, Calif; Henry A. Pitt, MD, Baltimore, Md | |
| ● In patients with acute cholecystitis with two to three predictive factors, antibiotics should be continued until culture data are available. | |
| The 'Plug' Repair of 1402 Recurrent Inguinal Hernias: 20-Year Experience | 265 |
| Alex G. Shulman, MD; Parviz K. Amid, MD; Irving L. Lichtenstein, MD, Los Angeles, Calif | |
| ● The "plug technique" to occlude the hernial defect without tension results in a low failure rate, minimal postoperative discomfort, and rapid rehabilitation. | |
| Invited Commentary: Lloyd M. Nyhus, MD, Chicago, Ill | |
| Adult Assault as a Mechanism of Burn Injury | 268 |
| Gary F. Purdue, MD, John L. Hunt, MD, Dallas, Tex | |
| ● Assault by burning in adults is often not appreciated as a cause of burn injury. | |
| CLINICAL NOTES | |
| Postoperative Chylous Ascites: Diagnosis and Treatment: A Series Report and Literature Review | 270 |
| C. John Ablan, MD; Fred N. Littooy, MD; Robert J. Freeark, MD, Maywood, Ill | |
| ● Early recognition and conservative management of chyle ascites is most often effective with peritoneovenous shunting reserved for treatment failures. | |
| Neuroleptic Malignant Syndrome in a Multiple Trauma Patient | 274 |
| Richard C. Boorse, MD, Michael Rhodes, MD, Allentown, Pa | |
| REGULAR DEPARTMENTS | |
| Books | 276 |
| Calendar of Events | 276 |
| Instructions for Authors | 177 |
| Classified Advertising | 277 |
| Index to Advertisers | 136 |
| CORRECTION | |
| Improving Diagnosis of Appendicitis: Early Autologous Leukocyte Scanning | 260 |
| Anne R. DeLaney, MD; Carol A. Raviola, MD; Paul N. Weber, MD; Paul T. McDonald, MD; Daniel A. Navarro, MD; Ivan Jasko, Oakland, Calif | |