

Copyright © 1990 by the AMERICAN MEDICAL ASSOCIATION

Official Publication of the Association of Veterans Administration Surgeons, New England Surgical Society, Pacific Coast Surgical Association, Society of Surgical Oncology, Surgical Infection Society, Western Surgical Association.

COMMENTARY

- Sabbatical Leave for Surgeons** 297
Stanley R. Friesen, MD, PhD, Kansas City, Kan

SPECIAL ARTICLE

- Mammographic Second Opinions Prior to Biopsy of Nonpalpable Breast Lesions** 298
Ferris M. Hall, MD, Boston, Mass
Invited Commentary: Alexander J. Walt, MD, Detroit, Mich

PAPERS READ BEFORE THE ANNUAL MEETING OF THE SOCIETY OF SURGICAL**ONCOLOGY, SAN FRANCISCO, CALIF, MAY 21 TO MAY 24, 1989—PART II**

- Carcinoembryonic Antigen Enhances Metastatic Potential of Human Colorectal Carcinoma** 300
Richard B. Hostetter, MD; Debora E. Campbell; Kefung Chi, MD; Susan Kerckhoff; Karen R. Cleary, MD;
Steve Ullrich, PhD, Houston, Tex; Peter Thomas, PhD, Boston, Mass; J. Milburn Jessup, MD, Houston, Tex
• Carcinoembryonic antigen enhanced experimental metastasis in a low metastatic human colorectal carcinoma (KM-12c) but did not alter the metastatic potential or high metastatic or nonmetastatic lines.
- Seroma Prevention Using Fibrin Glue in a Rat Mastectomy Model** 305
William H. Lindsey; Thomas M. Masterson, MD; William D. Spotnitz, MD;
Morton C. Wilhelm, MD; Raymond F. Morgan, MD, Charlottesville, Va
• Topical fibrin glue is statistically significant in decreasing postmastectomy seroma formation in the Sprague-Dawley rat.
- Tumor Immunization: Improved Results After Vaccine Modified With Recombinant Interferon Gamma** 308
Robert K. Sigal, MD; Michael D. Lieberman, MD; John V. Reynolds, FRCS; Noel Williams, FRCS;
Moritz M. Ziegler, MD; John M. Daly, MD, Philadelphia, Pa
• Modification of tumor with recombinant interferon gamma before introduction as a vaccine may improve that vaccine's protective capability.
- Patterns of Failure in Anorectal Melanoma: A Guide to Surgical Therapy** 313
Merrick Ross, MD; Christopher Pezzi, MD; Thomas Pezzi; Dennis Meurer, MD;
Robert Hickey, MD; Charles Balch, MD, Houston, Tex
• Local excision is recommended when technically feasible since these patients eventually succumb to metastasis regardless of surgical therapy.
- Isolated Limb Perfusion for Localized Melanoma of the Extremity: A Matched Comparison of Wide Local Excision With Isolated Limb Perfusion and Wide Local Excision Alone** 317
Michael J. Edwards, MD, Houston, Tex; Seng-Jaw Soong, PhD, Birmingham, Ala; Arthur W. Boddie, MD;
Charles M. Balch, MD; Charles M. McBride, MD, Houston, Tex
• Isolated limb perfusion with melphalan may improve survival in selected patients with malignant melanoma and justifies continued prospective trials.
- Prognosis of Thick Cutaneous Melanoma of the Trunk and Extremity** 322
Daniel Coit, MD; Paul Sauven, MD, FRCS; Murray Brennan, MD, New York, NY
• Half of the patients with extremity and truncal TCM present with locoregional disease warranting aggressive surgical approach; with negative nodes the 5-year survival is superior to previous reports.
- Stage B Prostate Adenocarcinoma: Flow Cytometric Nuclear DNA Ploidy Analysis** 327
B. T. Montgomery, MD; Ofer Nativ, MD; Michael L. Blute, MD; George M. Farrow, MD; Robert P. Myers, MD;
Horst Zincke, MD; Terry M. Therneau, PhD; Michael M. Lieber, MD, Rochester, Minn
• Nuclear DNA content has prognostic significance for patients with prostatic adenocarcinoma.
Invited Commentary: Donald G. Skinner, MD, Los Angeles, Calif
- Ablation of Abnormal Energy Expenditure by Curative Tumor Resection** 337
James D. Luketich, MD; James L. Mullen, MD; Irene D. Feurer, MSEd;
Jonathan Sternlieb, MD; Robert C. Fried, MD, Philadelphia, Pa
• Abnormal energy expenditure in patients with cancer cannot be attributed solely to abnormal host body composition.

The ARCHIVES OF SURGERY (ISSN 0004-0010) is published monthly by the American Medical Association, 535 N Dearborn St, Chicago, IL 60610, and is an official publication of the Association. Second-class postage paid at Chicago, IL 60610, and at additional mailing office.

SUBSCRIPTION RATES—The subscription rates for the ARCHIVES OF SURGERY are as follows: \$56 for one year, \$97 for two years in the United States and US possessions; all other countries, one year, \$66; two years, \$117 for surface delivery. For expedited air delivery to most countries, add \$15 surcharge for one year subscription, \$30 for two years. (Rates for subscriptions for delivery to Japan or South Korea are available through exclusive agents—contact the publisher.) **Special rates for residents and medical students in the United States and US Possessions are available.** Address inquiries to the American Medical Association, Circulation

and Fulfillment Division, 535 N Dearborn St, Chicago, IL 60610. Phone: (312)280-7168.

CHANGE OF ADDRESS—POSTMASTER, send all address changes to the ARCHIVES OF SURGERY, Attention: Subscription Department, 535 N Dearborn St, Chicago, IL 60610. Notification of address change must be made at least six weeks in advance, include both old and new addresses, a recent mailing label, and your new zip code.

Advertising Principles: Each advertisement in this issue has been reviewed and complies with the principles governing advertising in AMA scientific publications. A copy of these principles is available on request. The appearance of advertising in AMA publications is not an AMA guarantee or endorsement of the product or the claims made for the product by the manufacturer.

CONTENTS—Continued

Induction of the Expression of Differentiation-Related Antigens on Human Colon Carcinoma Cells by Stimulating Protein Kinase C	344
Paul L. Baron, MD; Michael J. Koretz, MD; Richard A. Carchman, PhD; James M. Collins, PhD; Anthony S. Tokarz, MS; George A. Parker, MD, Richmond, Va	
● PMA causes changes in structure, cell growth kinetics, and antigen expression, consistent with induction of differentiation of cell line SW 48.	
Inhibition of Sialic Acid Incorporation Prevents Hepatic Metastases	351
Hans E. Wagner, MD; Peter Thomas, PhD; Barbara C. Wolf, MD; Alan Rapoza; Glenn Steele, Jr, MD, PhD, Boston, Mass	
● Depletion of sialic acid from cell surface glycoconjugates reduces the incidence of hepatic metastases from human colorectal cancer.	
Combination Regional Therapy for Extremity Sarcoma: A Tricenter Study	355
Harold J. Wanebo, MD, Providence, RI; Walley J. Temple, MD, Calgary, Canada; Martin B. Popp, MD, Charles E. Douvill, MD, Cincinnati, Ohio; Michael Yablonski, Providence, RI	
● Combined therapy for extremity sarcomas resulted in excellent local control, good function in patients undergoing limb-salvage surgery, and reasonable long-term survival.	
Costs, Quality, and the Volume of Surgical Oncology Procedures	360
Eric Munoz, MD, MBA; Katherine Mulloy, AS; Jonathan Goldstein, MPA; Noel Tenenbaum, MD; Leslie Wise, MD, New Hyde Park, NY	
● Hospital cost and perhaps outcome may be volume related: the diagnosis related group prospective payment system may provide disincentives for low-volume surgeons.	
Invited Commentary: Joseph A. Ignatius, MD, Los Altos Hills, Calif	
Segmental Mastectomy Without Radiotherapy for T1 and Small T2 Breast Carcinomas	364
Frederick L. Moffat MD, FRCSC; Alfred S. Ketcham, MD; David S. Robinson, MD; Adrian Legaspi, MD; Latifa Ghandur-Mnaymneh, MD; Susan Hilsenbeck, MS, Miami, Fla	
● Every patient treated with segmental mastectomy may not require radiotherapy; studies are needed to define these exceptions.	
Intraoperative Radiation Therapy and Hyperthermia: Morbidity and Mortality From This Combined Treatment Modality for Unresectable Intra-abdominal Carcinomas	370
Thomas A. Colacchio, MD; Christopher Coughlin, MD; James Taylor, MD; Evan Douple, PhD; Thomas Ryan, MS; Robert W. Crichlow, MD, Hanover, NH	
● A mortality of 11% and a morbidity of 58% reveal a continued need for improved technology in the use of intraoperative hyperthermia for large, complexly shaped intra-abdominal tumors.	
Axillary Lymphadenectomy: A Prospective, Randomized Trial of 13 Factors Influencing Drainage, Including Early or Delayed Arm Mobilization	378
Jeanne A. Petrek, MD; Margaret M. Peters, RN; Subhadra Nori, MD; Cynthia Knauer, RN; David W. Kinne, MD; Andre Rogatko, PhD, New York, NY	
● Early vs delayed time of exercise initiation following axillary lymphadenectomy had no effect on the amount or duration of drainage in inpatients or outpatients.	
ORIGINAL ARTICLES	
Comparison of Magnetic Resonance Imaging and Computed Tomography in the Preoperative Staging of Rectal Cancer	385
Claude Guinet, MD, PhD; Jean-Noël Buy, MD; Michel A. Ghossain, MD; Alain Sézeur, MD; Alain Mallet, MD; Jean-Michel Bigot, MD; Dominique Vadrot, MD; Jean Ecoiffier, MD, Paris, France	
● No significant discordance was observed between CT and MRI in preoperative staging of rectal cancer.	
Achievements of Physical Therapy in Patients After Modified Radical Mastectomy Compared With Quadrantectomy, Axillary Dissection, and Radiation for Carcinoma of the Breast	389
Haim Gutman, MD; Tali Kersz; Tali Barzilai; Menashe Haddad, MD; Raphael Reiss, MD, Petah Tiqva, Israel	
● Patients undergoing quadrantectomy, axillary dissection, and radiation therapy suffer less pain and require less intensive physical therapy.	
Growth Delay in Postburn Pediatric Patients	392
Randi L. Rutan, RN, BSN, David N. Herndon, MD, Galveston, Tex	
● The exact cause of growth delay following severe thermal injury in the pediatric population is unknown.	
Insulin Protects Against Muscle Proteolysis Induced by Septic Plasma	396
Linda Mitchell, MD, Lawrence W. Norton, MD, Denver, Colo	
● Proteolysis can be decreased in septic patients by the administration of insulin.	
Surgery for Nonpalpable Breast Lesions	399
Angelos E. Papatestas, MD; Dorit Hermann; George Hermann, MD; Constantinos Tsevdos; Gerson Lesnick, MD, New York, NY	
● Spot localization is more appropriate than monitoring to achieve earlier recognition of and therapy for occult neoplasms.	
BASIC SCIENCE REVIEW	
The Role of Intestinal Barrier Failure and Bacterial Translocation in the Development of Systemic Infection and Multiple Organ Failure	403
Edwin A. Deitch, MD, FACS, Shreveport, La	
CLINICAL NOTES	
The Retained Surgical Sponge Following Intra-abdominal Surgery: A Continuing Problem	405
William Rappaport, MD, Kevin Haynes, MD, Tucson, Ariz	
● The retained foreign body following laparotomy is a continuing problem with variable presentations.	
Intracellular Lipid Droplets in Functioning Transitional Parathyroid Oxyphil Adenomas: A Caveat	410
Harlan D. Alpern, MD, Cooperstown, NY; Sanford I. Roth, MD, Chicago, Ill; John E. Olson, MD, Cooperstown, NY	
REGULAR DEPARTMENTS	
Books	412
Calendar of Events	412
Instructions for Authors	295
Classified Advertising	413
Index to Advertisers	296